

Appendix A. Search Strategy

RTI MDD Search Strategy

Pubmed 02.05.2014

Search	Query	Items found
#1	Search ("Bupropion"[Mesh] OR "Bupropion"[tiab] OR 34911-55-2[rn])	3570
#2	Search ("Citalopram"[Mesh] OR "Citalopram"[tiab] OR 59729-33-8[rn])	5061
#3	Search ("Escitalopram"[tiab] OR 128196-01-0[rn])	4031
#4	Search ("O-desmethylvenlafaxine" [Supplementary Concept] OR Desvenlafaxine[tiab] OR 93413-62-8[rn])	234
#5	Search ("Fluoxetine"[Mesh] OR "Fluoxetine"[tiab] OR 54910-89-3[rn])	10911
#6	Search ("Fluvoxamine"[Mesh] OR "Fluvoxamine"[tiab] OR 54739-18-3[rn])	2551
#7	Search (("milnacipran"[Supplementary Concept] OR "Levomilnacipran"[tiab] OR 96847-54-0[rn]))	346
#8	Search ("mirtazapine"[Supplementary Concept] OR "mirtazapine"[tiab] OR 85650-52-8[rn])	1574
#9	Search ("nefazodone"[Supplementary Concept] OR "nefazodone"[tiab] OR 82752-99-6[rn])	706
#10	Search ("Paroxetine"[Mesh] OR "Paroxetine"[tiab] OR 61869-08-7[rn])	5220
#11	Search ("Sertraline"[Mesh] OR "Sertraline"[tiab] OR 79617-96-2[rn])	3732
#12	Search ("Trazodone"[Mesh] OR "Trazodone"[tiab] OR 19794-93-5[rn])	1685
#13	Search ("venlafaxine"[Supplementary Concept] OR "venlafaxine"[tiab] OR 93413-69-5[rn])	3124
#14	Search ("vilazodone"[Supplementary Concept] OR "vilazodone"[tiab] OR 163521-12-8[rn])	65
#15	Search ("vortioxetine"[Supplementary Concept] OR "vortioxetine"[tiab] OR 508233-74-7[rn])	50
#16	Search ("Antidepressive Agents, Second-Generation"[Mesh] OR "Antidepressive Agents, Second-Generation"[Pharmacological Action])	57579
#17	Search (#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16)	67046
#18	Search ("Psychotherapy"[Mesh] OR psychotherap*[tiab])	156412
#19	Search (Acceptance and Commitment Therap*[tiab] OR Cognitive Therap*[tiab] OR Cognitive behavioral Therap*[tiab] OR interpersonal therap*[tiab] OR psychodynamic therap*[tiab] OR behavioral therap*[tiab])	8028
#20	Search (#18 OR #19)	158665
#21	Search "Hypericum"[Mesh] OR "Hypericum"[tiab] OR "St. Johns Wort"[tiab] OR "Saint Johns Wort"[tiab] OR "St. John's Wort"[tiab] OR "Saint John's Wort"[tiab] OR LI160[tiab] OR LI160[tiab] OR WS5572[tiab] OR WS5573[tiab] OR LoHyp-57[tiab]	2591
#22	Search "s adenosyl l methionine"[tiab] OR "s adenosylmethionine"[tiab] OR "S-Adenosylmethionine"[Mesh]	8814
#23	Search "Fatty Acids, Omega-3"[Mesh] OR (omega 3[tiab] AND fatty acid*[tiab]) OR fish oil[tiab] OR flax seed[tiab] OR borage seed[tiab] OR Borago[tiab] OR evening primrose[tiab] OR Oenothera[tiab] OR eicosapentaenoic acid[tiab] OR PUFA[tiab]	27732
#24	Search "Acupuncture"[Mesh] OR "Acupuncture Therapy"[Mesh] OR Acupuncture[tiab] OR Electroacupuncture[tiab]	20337
#25	Search "Yoga"[Mesh] OR yoga[tiab]	2387
#26	Search "Meditation"[Mesh] OR meditation[tiab] OR mindfulness[tiab]	4174
#27	Search ("Exercise"[Mesh] OR physical activit*[tiab] OR "physical exercise"[tiab])	159734
#28	Search (#22 OR #23 OR #24 OR #25 OR #26 OR #27)	221143
#29	Search ("Depressive Disorder, Major"[MeSH] OR "major depressive disorder"[tiab] OR "major depression"[tiab])	35489
#30	Search (#29 AND (#28 OR #20 OR #17))	7950
#31	Search (systematic*[tiab] AND (bibliographic*[tiab] OR literature[tiab] OR review[tiab] OR reviewed[tiab] OR reviews[tiab])) OR (comprehensive*[tiab] AND	241160

Search	Query	Items found
	(bibliographic*[tiab] OR literature[tiab])) OR "research synthesis"[tiab] OR "research integration"[tiab] OR meta-analy*[tiab] OR metaanaly*[tiab] OR "meta-analysis as topic"[mh] OR "Meta-Analysis"[pt] OR ("review"[tiab] AND ("rationale"[tiab] OR "evidence"[tiab]) AND review[pt]) OR "Systematic Review"[tiab] OR ("Review"[Publication Type] AND "systematic"[tiab])	
#32	Search ("Randomized Controlled Trial"[Publication Type] OR "Randomized Controlled Trials as Topic"[MeSH] OR "Randomized Controlled Trial"[tiab] OR "Single-Blind Method"[MeSH] OR "Double-Blind Method"[MeSH] OR "Random Allocation"[MeSH])	542163
#33	Search (("cohort studies"[MeSH] OR cohort stud*[tiab] OR cohort analy*[tiab] OR "Case-Control Studies"[Mesh] OR case control stud*[tiab] OR observational stud*[tiab] OR "observational study"[pt] OR ((longitudinal[tiab] OR retrospective[tiab] OR prospective[tiab]) AND (study[tiab] OR trial[tiab]))) AND ("Comparative Study"[pt] OR comparison[tiab] OR comparative[tiab]))	349594
#34	Search ("Controlled Clinical Trial"[pt] OR "Controlled Clinical Trials as Topic"[Mesh] OR controlled clinical trial*[tiab] OR controlled trial*[tiab] OR controlled stud*[tiab])	319105
#35	Search (#30 AND (#31 OR #32 OR #33 OR #34))	3642
#36	Search ("Animals"[Mesh] NOT "Humans"[Mesh])	3882887
#37	Search (#35 NOT #36)	3635
#38	Search ("Infant"[Mesh] OR "Child"[Mesh] OR "Adolescent"[Mesh]) NOT "Adult"[Mesh]	1490657
#39	Search (#37 NOT #38)	3398
#40	Search #39 AND 1990:2014[dp] AND (english[la] OR german[la] OR italian[la])	3231

Addendum duloxetine 07.05.2014

Search	Query	Items found
#1	Search ("duloxetine" [Supplementary Concept] OR duloxetine[tiab])	1677
#2	Search ("Depressive Disorder, Major"[MeSH] OR "major depressive disorder"[tiab] OR "major depression"[tiab])	35518
#3	Search (#1 AND #2)	430
#4	Search ((systematic*[tiab] AND (bibliographic*[tiab] OR literature[tiab] OR review[tiab] OR reviewed[tiab] OR reviews[tiab])) OR (comprehensive*[tiab] AND (bibliographic*[tiab] OR literature[tiab])) OR "research synthesis"[tiab] OR "research integration"[tiab] OR meta-analy*[tiab] OR metaanaly*[tiab] OR "meta-analysis as topic"[mh] OR "Meta-Analysis"[pt] OR ("review"[tiab] AND ("rationale"[tiab] OR "evidence"[tiab]) AND review[pt]) OR "Systematic Review"[tiab] OR ("Review"[Publication Type] AND "systematic"[tiab]))	241577
#5	Search (("Randomized Controlled Trial"[Publication Type] OR "Randomized Controlled Trials as Topic"[MeSH] OR "Randomized Controlled Trial"[tiab] OR "Single-Blind Method"[MeSH] OR "Double-Blind Method"[MeSH] OR "Random Allocation"[MeSH]))	542686
#6	Search (((("cohort studies"[MeSH] OR cohort stud*[tiab] OR cohort analy*[tiab] OR "Case-Control Studies"[Mesh] OR case control stud*[tiab] OR observational stud*[tiab] OR "observational study"[pt] OR ((longitudinal[tiab] OR retrospective[tiab] OR prospective[tiab]) AND (study[tiab] OR trial[tiab]))) AND ("Comparative Study"[pt] OR comparison[tiab] OR comparative[tiab])))	349872
#7	Search (((("Controlled Clinical Trial"[pt] OR "Controlled Clinical Trials as Topic"[Mesh] OR controlled clinical trial*[tiab] OR controlled trial*[tiab] OR controlled stud*[tiab])))	319351
#8	Search (#3 AND (#7 OR #6 OR #5 OR #4))	234
#9	Search (("Animals"[Mesh] NOT "Humans"[Mesh]))	3884483
#10	Search (#8 NOT #9)	234
#11	Search (((("Infant"[Mesh] OR "Child"[Mesh] OR "Adolescent"[Mesh]) NOT "Adult"[Mesh]))	1491426
#12	Search (#10 NOT #11)	234
#13	Search (#12 AND 1990:2014[dp] AND (english[la] OR german[la] OR italian[la]))	229

Cochrane Library 02.05.2014

ID	Search	Hits
#1	[mh Bupropion] or "Bupropion":ti,ab	934
#2	[mh Citalopram] or "Citalopram":ti,ab	1196
#3	Escitalopram:ti,ab	588
#4	Desvenlafaxine:ti,ab	71
#5	[mh Fluoxetine] or "Fluoxetine":ti,ab	2364
#6	[mh Fluvoxamine] or "Fluvoxamine":ti,ab	687
#7	Levomilnacipran:ti,ab	12
#8	mirtazapine:ti,ab	439
#9	nefazodone:ti,ab	194
#10	[mh Paroxetine] or "Paroxetine":ti,ab	1770
#11	[mh Sertraline] or "Sertraline":ti,ab	1350
#12	[mh Trazodone] or "Trazodone":ti,ab	335
#13	venlafaxine:ti,ab	989
#14	vilazodone:ti,ab	19
#15	vortioxetine:ti,ab	13
#16	[mh "Antidepressive Agents, Second-Generation"]	1230
#17	{or #1-#16}	9063
#18	[mh Psychotherapy] or psychotherap*:ti,ab	17066
#19	(Acceptance near/2 Commitment next Therap*):ti,ab or (Cognitive near/2 Therap*):ti,ab or ((interpersonal or psychodynamic or behavioral) next therap*):ti,ab	5015
#20	#18 or #19	19087
#21	[mh yoga] or yoga:ti,ab	688
#22	[mh meditation] or (meditation or mindfulness):ti,ab	1064
#23	[mh Acupuncture] or [mh "Acupuncture Therapy"] or (Acupuncture or Electroacupuncture):ti,ab	7130
#24	[mh Hypericum] or "Hypericum":ti,ab or (john* next wort):ti,ab or (LI160 or WS5572 or WS5573 or LoHyp-57):ti,ab	295
#25	("s adenosyl l methionine" or "s adenosylmethionine"):ti,ab or [mh S-Adenosylmethionine]	191
#26	[mh "fatty Acids, Omega-3"] or (omega-3 and fatty next acid*):ti,ab or ("fish oil" or "flax seed" or "borage seed" or Borago or "evening primrose" or Oenothera or "eicosapentaenoic acid" or PUFA):ti,ab	3661
#27	[mh Exercise] or (physical next (activit* or exercise)):ti,ab	18551
#28	{or #21-#27}	31146
#29	[mh "Depressive Disorder, Major"] or "major depressive disorder":ti,ab or (major next/1 depress*):ti,ab	6406
#30	#29 and (#17 or #20 or #28)	3544
#31	#30 Publication Date from 1990 to 2014	3460
#32	(([mh infant] or [mh child] or [mh adolescent]) not [mh adult])	88056
#33	#31 not #32	2867
#34	#31 and (adult or adults):ti,ab	393
#35	#33 or #34	2945
#36	[mh animals] not [mh humans]	5655
#37	#35 not #36 in Other Reviews, Trials, Methods Studies, Technology Assessments, Economic Evaluations and Cochrane Groups	2940
#38	(review:pt and systematic:ti,ab) or "systematic review"	37971
#39	meta-analysis:pt or (meta next analy*):ti,ab or metaanaly*:ti,ab or [mh Meta-Analysis] or [mh "Meta-Analysis as Topic"]	20924
#40	[mh "Randomized Controlled Trial"] or [mh "Randomized Controlled Trial as topic"] or "randomized controlled trial":pt or [mh "single-blind method"] or [mh "double-blind method"] or [mh "random allocation"] or (randomi?ed next controlled next "trial"):ti,ab	377221
#41	[mh "Controlled Clinical Trial"] or [mh "Controlled Clinical Trials as Topic"] or (controlled next/2 (trial or study)):ti,ab	149716
#42	(([mh "cohort studies"] or (cohort next stud*):ti,ab or [mh "case-control studies"] or (case-control next stud*):ti,ab or (observational next stud*):ti,ab or "observational study":pt or ((observational or longitudinal or retrospective) near/2 (study or trial)):ti,ab) and ("comparative study":pt or [mh "Comparative Study"] or comparison:ti,ab or	51743

ID	Search	Hits
	comparative:ti,ab)	
#43	{or #38-#42}	458686
#44	#37 and #43 in Other Reviews, Trials, Methods Studies, Technology Assessments, Economic Evaluations and Cochrane Groups	2000

Addendum duloxetine 07.05.2014

No.	Query	Results
#30	[mh "Depressive Disorder, Major"] or "major depressive disorder":ti,ab or (major next/1 depress*):ti,ab	6406
#31	#30 and "duloxetine":ti,ab	163
#32	#31 Publication Date from 1990 to 2014	163
#33	(([mh infant] or [mh child] or [mh adolescent]) not [mh adult])	88056
#34	#32 not #33	143
#35	#32 and (adult or adults):ti,ab	29
#36	#34 or #35	149
#37	[mh animals] not [mh humans]	5655
#38	#36 not #37 in Other Reviews, Trials, Methods Studies, Technology Assessments, Economic Evaluations and Cochrane Groups	149
#39	[mh "Randomized Controlled Trial"] or [mh "Randomized Controlled Trial as topic"] or "randomized controlled trial":pt or [mh "single-blind method"] or [mh "double-blind method"] or [mh "random allocation"] or (randomi?ed next controlled next "trial"):ti,ab	377221
#40	[mh "Controlled Clinical Trial"] or [mh "Controlled Clinical Trials as Topic"] or (controlled next/2 (trial or study)):ti,ab	149722
#41	#38 and (#39 or #40) in Trials	87
#42	#38 in Other Reviews	8
#43	#42 or #41	95

EMBASE 06.05.2014

No.	Query	Results
#1.1	'amfebutamone'/exp OR bupropion:tn,ab,ti OR '34911 55 2':rn	
#1.2	'citalopram'/exp OR citalopram:tn,ab,ti OR '59729 33 8':rn OR 'escitalopram'/exp OR escitalopram:tn,ab,ti OR '128196 01 0':rn	
#1.3	'desvenlafaxine'/exp OR desvenlafaxine:tn,ab,ti OR '93413 62 8':rn	
#1.4	'fluoxetine'/exp OR fluoxetine:tn,ab,ti OR '54910 89 3':rn	
#1.5	'fluvoxamine'/exp OR fluvoxamine:tn,ab,ti OR '54739 18 3':rn	
#1.6	'milnacipran'/exp OR levomilnacipran:tn,ab,ti OR '96847 54 0':rn	
#1.7	'mirtazapine'/exp OR mirtazapine:tn,ab,ti OR '85650 52 8':rn	
#1.8	'nefazodone'/exp OR nefazodone:tn,ab,ti OR '82752 99 6':rn	
#1.9	'paroxetine'/exp OR paroxetine:tn,ab,ti OR '61869 08 7':rn	
#1.10	'sertraline'/exp OR sertraline:tn,ab,ti OR '79617 96 2':rn	
#1.11	'trazodone'/exp OR trazodone:tn,ab,ti OR '19794 93 5':rn	
#1.12	'venlafaxine'/exp OR venlafaxine:tn,ab,ti OR '93413 69 5':rn	
#1.13	'vilazodone'/exp OR vilazodone:tn,ab,ti OR '163521 12 8':rn	
#1.14	'vortioxetine'/exp OR vortioxetine:tn,ab,ti OR '508233 74 7':rn	
#1.15	'antidepressant agent'/exp AND 'second generation':ab,ti	
#1.16	#1.1 OR #1.2 OR #1.3 OR #1.4 OR #1.5 OR #1.6 OR #1.7 OR #1.8 OR #1.9 OR #1.10 OR #1.11 OR #1.12 OR #1.13 OR #1.14 OR #1.15	87062
#1.17	'psychotherapy'/exp	187201
#1.18	((acceptance OR cognitive OR interpersonal OR psychodynamic OR behavioral) NEXT/3 (therapy OR therapies OR psychotherapy)):ab,ti	20374
#1.19	#1.17 OR #1.18 AND 'treatment outcome'/exp	25169
#1.20	'hypericum'/exp OR hypericum:ab,ti OR (john* NEXT/1 wort):ab,ti OR li160:ab,ti OR ws5572:ab,ti OR ws5573:ab,ti OR 'lohyp 57':ab,ti	4193
#1.21	's adenosylmethionine'/exp OR 's adenosylmethionine' OR 's adenosyl l methionine':ab,ti OR 's adenosylmethionine':ab,ti	10557

No.	Query	Results
#1.22	'omega 3 fatty acid'/exp OR ('omega 3':ab,ti AND acid*:ab,ti) OR 'fish oil':ab,ti OR 'flax seed':ab,ti OR 'borage seed':ab,ti OR borago:ab,ti OR 'evening primrose':ab,ti OR oenothera:ab,ti OR 'eicosapentaenoic acid':ab,ti OR pufa:ab,ti	34034
#1.23	'acupuncture'/exp OR acupuncture:ab,ti OR electroacupuncture:ab,ti	33369
#1.24	'yoga'/exp OR yoga:ab,ti	4138
#1.25	'meditation'/exp OR meditation:ab,ti OR mindfulness:ab,ti	6584
#1.26	'exercise'/exp	209232
#1.27	#1.20 OR #1.21 OR #1.22 OR #1.23 OR #1.24 OR #1.25 OR #1.26 AND 'treatment outcome'/exp	19807
#1.28	'major depression'/exp OR 'major depressive disorder':ab,ti OR (major NEXT/2 depress*):ab,ti	52013
#1.29	#1.28 AND (#1.27 OR #1.19 OR #1.16)	13000
#1.30	'systematic review'/exp OR 'meta analysis'/exp OR 'systematic review':ab,ti OR (meta NEXT/1 analy*):ab,ti OR metaanaly*:ab,ti OR (review:it AND systematic:ab,ti) OR (systematic:ab,ti AND (bibliographic:ab,ti OR literature:ab,ti OR review:ab,ti OR reviewed:ab,ti OR reviews:ab,ti)) OR 'research synthesis':ab,ti OR 'research integration':ab,ti OR (comprehensive*:ab,ti AND (bibliographic:ab,ti OR literature:ab,ti)) OR (review:it AND review:ab,ti AND (rationale:ab,ti OR evidence:ab,ti))	287380
#1.31	'randomized controlled trial'/exp OR (randomi?ed NEXT/1 'controlled trial'):ab,ti OR 'double blind procedure'/exp OR 'single blind procedure'/exp OR 'randomization'/exp OR 'random allocation':ab,ti OR (allocated NEXT/2 random*):ab,ti	441315
#1.32	'cohort analysis'/exp OR 'case control study'/exp OR 'observational study'/exp OR 'longitudinal study'/exp OR 'prospective study'/exp OR 'retrospective study'/exp OR (cohort NEXT/1 (stud* OR analy*)):ab,ti OR (observational OR 'case control') NEXT/1 stud* OR ((longitudinal OR retrospective OR prospective) NEXT/2 (trial OR study)):ab,ti AND ('comparative study'/exp OR comparative:ab,ti OR comparison:ab,ti)	149667
#1.33	'controlled clinical trial'/exp OR (controlled NEXT/2 (trial* OR stud*)):ab,ti	591400
#1.34	#1.29 AND (#1.30 OR #1.31 OR #1.32 OR #1.33)	
#1.35	'human'/exp	14753345
#1.36	#1.34 AND #1.35	4575
#1.37	'adult'/exp	5294678
#1.38	#1.36 AND #1.37	2399
#1.39	#1.38 AND [1990-2014]/py	2374
#1.40	#1.39 AND [english]/lim	2316
#1.41	#1.39 AND (german:la OR italian:la)	17
#1.42	#1.40 OR #1.41	2333
#1		2333
#2	#1 AND [embase]/lim	2232

Addendum duloxetine 07.05.2014

No.	Query	Results
#6	'randomized controlled trial'/exp OR (randomi?ed NEXT/1 'controlled trial'):ab,ti OR 'double blind procedure'/exp OR 'single blind procedure'/exp OR 'randomization'/exp OR 'random allocation':ab,ti OR (allocated NEXT/2 random*):ab,ti	441315
#7	'cohort analysis'/exp OR 'case control study'/exp OR 'observational study'/exp OR 'longitudinal study'/exp OR 'prospective study'/exp OR 'retrospective study'/exp OR (cohort NEXT/1 (stud* OR analy*)):ab,ti OR (observational OR 'case control') NEXT/1 stud* OR ((longitudinal OR retrospective OR prospective) NEXT/2 (trial OR study)):ab,ti AND ('comparative study'/exp OR comparative:ab,ti OR comparison:ab,ti)	149667
#8	'controlled clinical trial'/exp OR (controlled NEXT/2 (trial* OR stud*)):ab,ti	591400
#9	#5 OR #6 OR #7 OR #8	1017694
#10	#4 AND #9	582
#11	#10 AND 'human'/exp AND 'adult'/exp	210
#12	#11 AND [1990-2014]/py AND [embase]/lim	205
#13	#12 AND ([english]/lim OR german:la OR italian:la)	205

CINAHL (via Ebsco) 02.05.2014

#	Query	Results
S1	(MH "Bupropion") OR "Bupropion"	1,072
S2	(MH "Citalopram") OR "Citalopram"	644
S3	"Escitalopram"	202
S4	(MH "Desvenlafaxine Succinate") OR TX Desvenlafaxine	49
S5	(MH "Fluoxetine+") OR "Fluoxetine"	1,144
S6	(MH "Fluvoxamine Maleate") OR "Fluvoxamine"	145
S7	"Levomilnacipran"	5
S8	(MH "Mirtazapine") OR "mirtazapine"	255
S9	(MH "Nefazodone") OR "nefazodone"	68
S10	(MH "Paroxetine") OR "Paroxetine"	732
S11	(MH "Sertraline Hydrochloride") OR "Sertraline"	643
S12	(MH "Trazodone") OR "Trazodone"	168
S13	(MH "Venlafaxine+") OR "venlafaxine"	605
S14	"vilazodone"	11
S15	"vortioxetine"	5
S16	(MH "Antidepressive Agents, Second Generation+")	2,806
S17	S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7 OR S8 OR S9 OR S10 OR S11 OR S12 OR S13 OR S14 OR S15 OR S16	4,599
S18	(MH "Psychotherapy+") OR (TI psychotherap*) OR (AB psychotherap*)	86,97
S19	TI (("Acceptance and Commitment" OR cognitive OR behavioral OR psychodynamic OR interpersonal) N2 therap*) OR AB (("Acceptance and Commitment" OR cognitive OR behavioral OR psychodynamic OR interpersonal) N2 therap*)	4,172
S20	S18 OR S19	87,95
S21	(MH "Yoga+") OR (TI yoga) OR (AB yoga)	2,268
S22	(MH "Meditation") OR (TI (meditation OR mindfulness)) OR (AB (meditation OR mindfulness))	2,652
S23	(MH "St. John's Wort") OR "hypericum" OR (TI john* N2 wort) OR (AB john* N2 wort) OR (TI (LI160 OR WS5572 OR WS5573 OR LoHyp-57)) OR (AB (LI160 OR WS5572 OR WS5573 OR LoHyp-57))	932
S24	(MH "Fatty Acids, Omega-3+") OR (TI "omega 3" N1 fatty acid*) OR (AB "omega 3" N1 fatty acid*) OR (TI ("fish oil" OR "flax seed" OR "borage seed" OR Borago OR "evening primrose" OR Oenothera OR "eicosapentaenoic acid" OR PUFA)) OR (AB ("fish oil" OR "flax seed" OR "borage seed" OR Borago OR "evening primrose" OR Oenothera OR "eicosapentaenoic acid" OR PUFA))	4,996
S25	(MH "S-Adenosylmethionine") OR (TI ("s adenosyl l methionine" OR "s adenosylmethionine")) OR (AB ("s adenosyl l methionine" OR "s adenosylmethionine"))	204
S26	(MH "Acupuncture+") OR (TI (acupuncture OR electroacupuncture)) OR (AB (acupuncture OR electroacupuncture))	8,556
S27	(MH "Exercise+") OR TI (physical N1 (activit* OR exercise)) OR AB (physical N1 (activit* OR exercise))	65,646
S28	S21 OR S22 OR S23 OR S24 OR S25 OR S26 OR S27	83,353
S29	(MH "Depression+") AND (TX major N2 depress*)	3,393
S30	TI (major n2 depress* OR "major depressive disorder") OR AB (major n2 depress* OR "major depressive disorder")	3,998
S31	S29 OR S30	4,08
S32	S31 AND (S17 OR S20 OR S28)	1,115
S33	(MH "Animals") NOT (MH "Human")	24,505
S34	S32 NOT S33	1,114
S35	((MH "Infant") OR (MH "Child") OR (MH "Adolescence")) NOT (MH "Adult+")	221,397
S36	S34 NOT S35	1,012
S37	S36 AND (PY 1990-2014) AND (LA (english OR german Or italian))	1,003
S38	S37 NOT (PT (editorial OR letter OR commentary))	867

Addendum duloxetine 07.05.2014

#	Query	Results
S1	(MH "Duloxetine Hydrochloride") OR (TX Duloxetine)	378
S2	(MH "Depression+") AND (TX major N2 depress*)	3,397
S3	T1 (major n2 depress* OR "major depressive disorder") OR AB (major n2 depress* OR "major depressive disorder")	4
S4	S2 OR S3	4,082
S5	S4 AND S1	51
S6	(MH "Animals") NOT (MH "Human")	24,582
S7	S5 NOT S6	51
S8	((MH "Infant") OR (MH "Child") OR (MH "Adolescence")) NOT (MH "Adult+")	221,684
S9	S7 NOT S8	51
S10	S9 AND (PY 1990-2014) AND (LA (english OR german Or italian))	51
S11	S10 NOT (PT (editorial OR letter OR commentary))	50

AMED (via Ovid) 02.05.2014

#	Suchen	Ergebnisse
1	exp antidepressive agents/	272
2	Bupropion.mp.	15
3	Citalopram.mp.	9
4	Escitalopram.mp.	3
5	(Desvenlafaxine or O-desmethylvenlafaxine).mp.	0
6	Fluoxetine.mp.	50
7	Levomilnacipran.mp.	0
8	mirtazapine.mp.	6
9	(nefazodone or Paroxetine or Sertraline or Trazodone or venlafaxine or vilazodone or vortioxetine).mp.	69
10	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9	363
11	exp psychotherapy/	8542
12	((acceptance or cognitive or interpersonal or psychodynamic or behavioral) adj3 (therap\$ or psychotherap\$)).mp.	1395
13	11 or 12	8895
14	exp fatty acids/ or exp fish oils/ or (omega 3 and acid*).mp. or (flax seed or borage seed or Borago or evening primrose or Oenothera or eicosapentaenoic acid or PUFA).mp.	640
15	exp hypericum/ or (hypericum or (john\$ adj1 wort)).mp. or (LI160 or WS5572 or WS5573 or LoHyp-57).mp.	415
16	(S-Adenosylmethionine or s adenosyl l methionine).mp.	21
17	exp acupuncture/ or exp electroacupuncture/ or (acupuncture or electroacupuncture).mp.	9182
18	exp meditation/ or (meditation or mindfulness).mp.	658
19	exp Yoga/ or yoga.mp.	501
20	(physical adj1 (activit* or exercise)).mp. or exp Exercise/	9857
21	14 or 15 or 16 or 17 or 18 or 19 or 20	20984
22	exp depressive disorder/	890
23	((major adj2 depress\$) or major depressive disorder).mp.	352
24	22 or 23	1126
25	10 or 13 or 21	29843
26	24 and 25	283
27	26	283
28	limit 27 to yr="1990 -Current"	282
29	(exp infant/ or exp child/ or exp adolescent/) not exp adult/	15352
30	28 not 29	269
31	28 and (adult or adults).ti,ab.	39
32	30 or 31	271

Addendum duloxetine 07.05.2014

#	Suchen	Ergebnisse
1	duloxetine.mp.	22
2	exp depressive disorder/	894
3	((major adj2 depress\$) or major depressive disorder).mp.	353
4	2 or 3	1131
5	1 and 4	5

PsycInfo (via Ebsco) 02.05.2014

#	Query	Results
S1	TX (Bupropion OR Citalopram OR Escitalopram OR O-desmethylvenlafaxine OR Desvenlafaxine OR Fluoxetine OR Fluvoxamine OR Levomilnacipran OR mirtazapine OR nefazodone OR Paroxetine OR Sertraline OR Trazodone OR venlafaxine OR vilazodone OR vortioxetine)	15,857
S2	DE "antidepressant drugs" AND TX (second generation)	144
S3	S1 OR S2	15,958
S4	(DE "acceptance and commitment therapy") or ((DE "cognitive therapy") or (DE "behavior therapy")) OR (DE psychotherapy)	64,196
S5	TI ((acceptance and commitment therap*) OR (cognitive N2 therap*) OR (behavior* therap) OR (interpersonal therap*) OR (psychodynamic therap*)) OR AB ((acceptance and commitment therap*) OR (cognitive N2 therap*) OR (behavior* therap) OR (interpersonal therap*) OR (psychodynamic therap*))	24,311
S6	S4 OR S5	78,648
S7	(DE acupuncture) OR (TX (acupuncture OR electroacupuncture))	1,717
S8	(DE meditation) OR (TX (meditation OR mindfulness))	8,986
S9	(DE hypericum perforatum) OR (TX (hypericum OR (john* N1 wort) OR LI160 OR WS5572 OR WS5573 LoHyp-57))	392
S10	(DE yoga) OR (TX yoga)	1,907
S11	TX ((omega-3 N1 fatty acid*) OR "fish oil" OR "flax seed" OR "borage seed" OR Borago OR "evening primrose" OR Oenothera OR "eicosapentaenoic acid" OR PUFA)	982
S12	TX ("s adenosyl l methionine" OR "s adenosylmethionine")	205
S13	DE exercise OR TI physical activit* OR AB physical activit*	30,086
S14	S7 OR S8 OR S9 OR S10 OR S11 OR S12 OR S13	43,124
S15	DE ("major depressive disorder" OR "Major Depression") OR TI ("major depressive disorder" OR (Major N2 Depress*)) OR AB ("major depressive disorder" OR (Major N2 Depress*))	94,026
S16	S15 AND (S3 OR S6 OR S14)	13,953
S17	TI (controlled N2 (trial OR study)) OR AB (controlled N2 (trial OR study))	36,244
S18	TI (randomi*ed controlled trial) OR AB (randomi*ed controlled trial) OR TI (random* N4 (trial OR study)) OR AB (random* N4 (trial OR study))	39,503
S19	TI ("double-blind" OR (random* assigned) OR "single-blind") OR AB ("double-blind" OR (random* assigned) OR "single-blind")	42,931
S20	TI (systematic N3 (bibliographic OR literature OR review# OR reviewed)) OR AB (systematic N3 (bibliographic OR literature OR review# OR reviewed)) OR (comprehensive N3 (bibliographic OR literature)) OR ((TI "research integration") OR (AB "research integration")) OR ((TI "research synthesis") OR (AB "research synthesis")) OR ((TI metaanaly* OR meta-analy*) OR (AB metaanaly* OR meta-analy*)) OR (MR ("systematic review" OR "meta analysis"))	31,825
S21	((MR "longitudinal study" OR "retrospective study" OR "prospective study") OR (DE "cohort analysis") OR TI ((cohort N1 (analy* OR stud*)) OR ((observational OR "case control") N1 stud*) OR ((longitudinal OR retrospective OR prospective) N2 (trial OR study))) OR AB ((cohort N1 (analy* OR stud*)) OR ((observational OR "case control") N1 stud*) OR ((longitudinal OR retrospective OR prospective) N2 (trial OR study)))) AND (TI (comparative OR comparison) OR AB (comparative OR comparison))	10,731
S22	(S17 OR S18 OR S19 OR S20 OR S21) NOT ((ZZ "comment/reply") OR (ZZ "editorial") OR (ZZ "letter"))	113,102
S23	S16 AND S22	3,595
S24	((ZP "animal")) not ((ZP "human"))	279,088

#	Query	Results
S25	S23 NOT S24	3,586
S26	((((ZG "childhood (birth-12 yrs)") or (ZG "infancy (2-23 mo)") or ((ZG "adolescence (13-17 yrs)")))) not ((ZG "adulthood (18 yrs & older)"))	396,672
S27	S25 NOT S26	3,4
S28	LA (english OR german OR italian)	3,441,047
S29	PY 1990-2014	2,451,294
S30	S27 AND S28 AND S29	3,172

Addendum duloxetine 07.05.2014

#	Query	Results
S1	DE ("major depressive disorder" OR "Major Depression") OR TI ("major depressive disorder" OR (Major N2 Depress*)) OR AB ("major depressive disorder" OR (Major N2 Depress*))	94,027
S2	S1 AND (TX duloxetine)	375
S3	TI (controlled N2 (trial OR study)) OR AB (controlled N2 (trial OR study))	36,244
S4	TI (randomi*ed controlled trial) OR AB (randomi*ed controlled trial) OR TI (random* N4 (trial OR study)) OR AB (random* N4 (trial OR study))	39,503
S5	TI ("double-blind" OR (random* assigned) OR "single-blind") OR AB ("double-blind" OR (random* assigned) OR "single-blind")	42,931
S6	TI (systematic N3 (bibliographic OR literature OR review# OR reviewed)) OR AB (systematic N3 (bibliographic OR literature OR review# OR reviewed)) OR (comprehensive N3 (bibliographic OR literature)) OR ((TI "research integration") OR (AB "research integration")) OR ((TI "research synthesis") Or (AB "research synthesis")) OR ((TI metaanaly* OR meta-analy*) OR (AB metaanaly* OR meta-analy*)) OR (MR ("systematic review" OR "meta analysis"))	31,825
S7	((MR "longitudinal study" OR "retrospective study" OR "prospective study") OR (DE "cohort analysis") OR TI ((cohort N1 (analy* OR stud*)) OR ((observational OR "case control") N1 stud*) OR ((longitudinal OR retrospective OR prospective) N2 (trial OR study))) OR AB ((cohort N1 (analy* OR stud*)) OR ((observational OR "case control") N1 stud*) OR ((longitudinal OR retrospective OR prospective) N2 (trial OR study)))) AND (TI (comparative OR comparison) OR AB (comparative OR comparison))	10,731
S8	(S3 OR S4 OR S5 OR S6 OR S7) NOT ((ZZ "comment/reply") OR (ZZ "editorial") OR (ZZ "letter"))	113,102
S9	S2 AND S8	161
S10	((ZP "animal")) not ((ZP "human"))	279,088
S11	S9 NOT S10	161
S12	((((ZG "childhood (birth-12 yrs)") or (ZG "infancy (2-23 mo)") or ((ZG "adolescence (13-17 yrs)")))) not ((ZG "adulthood (18 yrs & older)"))	396,672
S13	S11 NOT S12	161
S14	LA (english OR german OR italian)	3,441,047
S15	PY 1990-2014	2,451,294
S16	S13 AND S14 AND S15	154

Grey Literature Search

ClinicalTrials.gov 04.06.2014

41 studies found for:

("major depressive disorder" OR "major depression") AND Bupropion Adult, Senior Phase 2, 3, 4

170 studies found for:

("major depressive disorder" OR "major depression") AND Citalopram Adult, Senior Phase 2, 3, 4

170 studies found for:

("major depressive disorder" OR "major depression") AND Escitalopram Adult, Senior Phase 2, 3, 4

35 studies found for:

("major depressive disorder" OR "major depression") AND Desvenlafaxine Adult, Senior Phase 2, 3, 4

45 studies found for:

("major depressive disorder" OR "major depression") AND Fluoxetine Adult, Senior Phase 2, 3, 4

6 studies found for:

("major depressive disorder" OR "major depression") AND Fluvoxamine Adult, Senior Phase 2, 3, 4

7 studies found for:

("major depressive disorder" OR "major depression") AND Levomilnacipran Adult, Senior Phase 2, 3, 4

21 studies found for:

("major depressive disorder" OR "major depression") AND mirtazapine Adult, Senior Phase 2, 3, 4

Found no studies with search of: ("major depressive disorder" OR "major depression") AND nefazodone Adult, Senior Phase 2, 3, 4

61 studies found for:

("major depressive disorder" OR "major depression") AND Paroxetine Adult, Senior Phase 2, 3, 4

66 studies found for:

("major depressive disorder" OR "major depression") AND Sertraline Adult, Senior Phase 2, 3, 4

4 studies found for:

("major depressive disorder" OR "major depression") AND Trazodone Adult, Senior Phase 2, 3, 4

66 studies found for:

("major depressive disorder" OR "major depression") AND venlafaxine Adult, Senior Phase 2, 3, 4

13 studies found for:

("major depressive disorder" OR "major depression") AND vilazodone Adult, Senior Phase 2, 3, 4

24 studies found for:

("major depressive disorder" OR "major depression") AND vortioxetine Adult, Senior Phase 2, 3, 4

74 studies found for:

("major depressive disorder" OR "major depression") AND duloxetine Adult, Senior Phase 2, 3, 4

Document: GreyLiterature.enl

ICTRP 04.06.2014

342 records for 243 trials found for: major depress* AND antidepress*

Document: ICTRP-040614.xlsx

Drugs@FDA 02.06.2014

Levomilnacipran:

http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.Set_Current_Drug&Applicant=FOREST%20LABS%20INC&ProductMktStatus=1&goto=Search.DrugDetails

Vilazodone:

http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.Set_Current_Drug&Applicant=FOREST%20LABS%20INC&ProductMktStatus=1&goto=Search.DrugDetails

Vortioxetine:

http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.Set_Current_Drug&Applicant=TAKEDA%20PHARMS%20USA&ProductMktStatus=1&goto=Search.DrugDetails

Folder: FDA

European Medicines Agency 02.06.2014

Levomilnacipran: 0

Vilazodone: 0

Vortioxetine:

http://www.ema.europa.eu:80/ema/index.jsp?curl=pages/medicines/human/medicines/002717/human_med_001714.jsp&mid=WC0b01ac058001d124

Folder: EMA

National Institute of Mental Health website 11.06.2014

Search Terms: "major depression", "major depressive disorder"

Folder: NIMH

American Psychological Association 11.06.2014

Search terms, "major depressive disorder", major depression"

Folder: APA

Scopus 16.06.2014

(TITLE-ABS-KEY(({major depressive disorder} OR {major depression}) OR KEY(({disorder, major depressive}))) AND (TITLE-ABS-KEY((bupropion OR citalopram OR escitalopram OR desvenlafaxine OR fluoxetine OR fluvoxamine OR levomilnacipran OR mirtazapine OR nefazodone OR paroxetine OR sertraline OR trazodone OR venlafaxine OR vilazodone OR vortioxetine OR duloxetine) OR ("Acceptance and Commitment Therapy" OR "Cognitive Therapy" OR "Cognitive behavioral Therapy" OR "interpersonal therapy" OR "psychodynamic therapy" OR "behavioral therapy") OR (hypericum OR "St. Johns Wort" OR "Saint Johns Wort" OR "St. John's Wort" OR "Saint John's Wort") OR ("s adenosyl l methionine" OR "S-Adenosylmethionine") OR ("omega 3") OR (acupuncture OR electroacupuncture) OR (yoga OR meditation OR mindfulness) OR ("physical activity" OR "physical exercise")) OR KEY(psychotherapy)) AND (TITLE-ABS-KEY(adult*)) AND (DOCTYPE(cp))

162 document results

Document: GreyLiterature.enl

Web of Science Conference Proceedings Citation Index- Science 16.06 2014

Set	Results	Search
		Indexes=CPCI-S Timespan=All years
# 1	3,944	TOPIC: ("major depressive disorder" OR "major depression")
		Indexes=CPCI-S Timespan=All years
# 2	4,014	TOPIC: (bupropion OR citalopram OR escitalopram OR desvenlafaxine OR fluoxetine OR fluvoxamine OR levomilnacipran OR mirtazapine OR nefazodone OR paroxetine OR sertraline OR trazodone OR venlafaxine OR vilazodone OR vortioxetine OR duloxetine)
		Indexes=CPCI-S Timespan=1990-2014
# 3	628	TOPIC: ("Acceptance and Commitment Therapy" OR "Cognitive Therapy" OR "Cognitive behavioral Therapy" OR "interpersonal therapy" OR "psychodynamic therapy" OR "behavioral therapy")
		Indexes=CPCI-S Timespan=1990-2014
# 4	404	TOPIC: (hypericum OR "St. Johns Wort" OR "Saint Johns Wort" OR "St. John's Wort" OR "Saint John's Wort")
		Indexes=CPCI-S Timespan=1990-2014
# 5	456	TOPIC: ("s adenosyl l methionine" OR "S-Adenosylmethionine")
		Indexes=CPCI-S Timespan=1990-2014
# 6	1,749	TOPIC: ("omega 3")
		Indexes=CPCI-S Timespan=1990-2014
# 7	1,068	TOPIC: (acupuncture OR electroacupuncture)
		Indexes=CPCI-S Timespan=1990-2014
# 8	517	TOPIC: (yoga OR meditation OR mindfulness)
		Indexes=CPCI-S Timespan=1990-2014
# 9	6,836	TOPIC: ("physical activity" OR "physical exercise")
		Indexes=CPCI-S Timespan=1990-2014
# 10	15,492	#9 OR #8 OR #7 OR #6 OR #5 OR #4 OR #3 OR #2
		Indexes=CPCI-S Timespan=1990-2014
# 11	874	#10 AND #1
		Indexes=CPCI-S Timespan=1990-2014
		Refined by: TOPIC: (adult*)
# 12	55	#10 AND #1

Document: GreyLiterature.enl

Appendix B. Cochrane Depression, Anxiety and Neurosis Group (CCDAN) Topic List: Intervention – Psychological Therapies¹

- **Behavior therapy / behavior modification**
 - Activity scheduling
 - Assertiveness training [CINAHL]
 - Aversion therapy [APA]
 - Covert sensitization [APA]
 - Behavior contracting [CINAHL]
 - Behavior modification
 - Biofeedback, psychology [MeSH]
 - Feedback, sensory [MeSH]
 - Contingency management [CINAHL]
 - Conversion therapy [APA]
 - Distraction therapy
 - Exposure therapy (APA)
 - Abreaction therapy
 - Sensitivity training
 - Systematic desensitization therapy (APA)
 - Eye movement desensitization reprocessing [MeSH]
 - Implosive therapy [APA, MeSH]
 - Pleasant events
 - Psychoeducation
 - Problem-focused
 - Reciprocal inhibition therapy (APA)
 - Relaxation techniques [CINAHL]
 - Autogenic training
 - Distraction [CINAHL]
 - Guided imagery [CINAHL]
 - Response cost (APA)
 - Sleep phase chronotherapy [MeSH]
 - Social skills training
 - Social effectiveness
- **Cognitive behavioral therapy [APA]**
 - Problem solving
 - Rational emotive therapy
 - Reality therapy
 - Restructuring
 - Role play
 - Schemas

¹ Cochrane Depression, Anxiety, and Neurosis Group. CCDAN Topic List: Intervention - Psychological therapies. 2013
http://ccdan.cochrane.org/sites/ccdan.cochrane.org/files/uploads/CCDAN%20topics%20list_psychological%20therapies%20for%20website.pdf. Accessed October 17, 2014.

- Self-control
- Stress management
- **Third wave cognitive behavioral therapies**
 - Acceptance and commitment therapy (ACT)
 - Behavioral activation
 - Cognitive behavioral analysis system of psychotherapy (CBASP)
 - Compassion-focused
 - Dialectical behavior therapy (DBT)
 - Diffusion
 - Functional analytic psychotherapy (FAP)
 - Metacognitive therapy
 - Mind training
 - Mindfulness
- **Psychodynamic therapies**
 - Brief psychotherapy
 - Countertransference
 - Freudian
 - Group therapy
 - Balint group therapy
 - Insight oriented therapy
 - Jungian
 - Kleinian
 - Object relations
 - Person centred therapy, client-centred therapy
 - Psychoanalytic therapy
 - Alderian therapy
 - Dream analysis
 - Free association
 - Self analysis
 - Short-term psychotherapy
 - Transference
- **Humanistic therapies**
 - Existential therapy
 - Experiential therapy
 - Process-experiential
 - Gestalt therapy
 - Expressive therapy
 - Griefwork
 - Rogerian
 - Non-directive therapy
 - Supportive therapy
 - Transactional analysis
- **Integrative therapies**
 - Cognitive analytical therapy
 - Counselling
 - Eclectic therapy

- Interpersonal therapy
 - Psychodynamic interpersonal therapy
- Multimodal
- Transtheoretical
- **Systemic therapies**
 - Conjoint therapy
 - Couples, marital or relationship therapy
 - Emotion focused therapy
 - Family therapy
 - Integrative behavioral couple therapy (IBCT)
 - Narrative therapy
 - Personal construct
 - Socioenvironmental therapy
 - Milieu therapy
 - Therapeutic community
 - Solution focused brief therapy
- **Other psychologically-oriented interventions**
 - Acting out
 - Age regression therapy
 - Art therapy
 - Bibliotherapy
 - Catharsis
 - Colour therapy
 - Crisis intervention
 - Dance therapy
 - Drama therapy
 - Emotional freedom techniqueso hypnotherapy
 - Autosuggestion
 - Neuro-linguistic programming (NLP)
 - Persuasion
 - Meditation [CINAHL]
 - Morita therapy
 - Music therapy
 - Play therapy
 - Primal therapy
 - Psychodrama
 - Reminiscence therapy
 - Sex therapy

Appendix C. Studies Excluded at the Full-Text Review Stage

Exclude Code	Exclude Reason
X1	Ineligible publication type
X2	Ineligible population(s)
X3	Ineligible or no intervention(s)
X4	Ineligible study design
X5	Ineligible or no comparison(s)
X6	Ineligible outcome(s)
X7	Does not answer a Key Question of the review
X8	Mixed treatment comparisons
X9	Systematic review without relevant meta-analysis
X10	Abstract only available

- | | |
|--|---|
| <p>1. St. John's wort ineffective for major depression. J Psychosoc Nurs Ment Health Serv. 2001;39(7):9-. Exclusion Code: X1</p> <p>2. Research notebook. Study shows St. John's wort ineffective for major depression. FDA Consum. 2002;36(3):8-. Exclusion Code: X1</p> <p>3. Treating late-life depression: pharmacotherapy or psychotherapy? Brown University Geriatric Psychopharmacology Update. 2006;10(12):3-4. Exclusion Code: X1</p> <p>4. Paroxetine plus psychotherapy for major depression in the elderly. Brown University Psychopharmacology Update. 2006;17(8):4-5. Exclusion Code: X1</p> <p>5. Paroxetine found to maintain quality of life in elderly patients with depression. Brown University Geriatric Psychopharmacology Update. 2007;11(11):1. Exclusion Code: X1</p> <p>6. Augmenting standard antidepressant treatment may help recovery in elderly. Brown University Geriatric Psychopharmacology Update. 2007;11(8):1. Exclusion Code: X1</p> <p>7. Cost effectiveness of therapy and fluoxetine for MDD in Romania... major depressive disorder. Brown University Psychopharmacology Update. 2009;20(4):4-. Exclusion Code: X1</p> <p>8. Better response and remission with mirtazapine for MD... major depression. Brown University Psychopharmacology Update. 2010;21(10):3-4. PMID: 2010780944. Language: English. Entry Date: 20101008. Revision Date: 20140103. Publication Type: journal article. Journal Subset: Biomedical. Exclusion Code: X1</p> | <p>9. Omega-3 effective in treating major depression - but only in absence of anxiety. Canadian Nursing Home. 2010;21(3):26-. Exclusion Code: X1</p> <p>10. [Therapy of moderately severe depressions in daily practice: first patient care research study reinforces clinical data]. MMW Fortschr Med. 2011 Oct 13;153(41):38-9. PMID: 22046838. Exclusion Code: X6</p> <p>11. A Double-Blind, Paroxetine- and Placebo-Controlled Study of 50 mg/Day and 100 mg/Day of EB-1010 Among Outpatients With Major Depressive Disorder Who Have Responded Inadequately to Prior Selective Serotonin Reuptake Inhibitors (SSRIs) and Serotonin Norepinephrine Reuptake Inhibitors (SNRIs) (Triple Reuptake Inhibitor Anti-Depressant Effects - TRIADE Study) [NCT01318434]. Clinicaltrials.gov [www.clinicaltrials.gov]; 2011. Exclusion Code: X9</p> <p>12. . Mittelschwere und schwere depression: langzeitbehandlung mit hypericum-extrakt WS 5570 gleich wirksam wie paroxetin. Schweiz Z Ganzheits Medizin. 2008 May;20(4):198-9. PMID: 0110887. Exclusion Code: X1</p> <p>13. Abt KL. The effects of a group exercise intervention in the adjunctive treatment of depression. US: ProQuest Information & Learning; 2006. Exclusion Code: X2</p> <p>14. Agid O, Lerer B. Algorithm-based treatment of major depression in an outpatient clinic: Clinical correlates of response to a specific serotonin reuptake inhibitor and to triiodothyronine augmentation. Int J Neuropsychopharmacol. 2003;6(1):41-9. Exclusion Code: X5</p> |
|--|---|

15. Alexopoulos GS, Canuso CM, Gharabawi GM, et al. Placebo-controlled study of relapse prevention with risperidone augmentation in older patients with resistant depression. *Am J Geriatr Psychiatry*. 2008 Jan;16(1):21-30. PMID: 17928573. Exclusion Code: X5
16. Alper BS. Evidence-based medicine. St. John's wort may be as effective as standard antidepressants--and more tolerable--for major depression. *Clinical Advisor for Nurse Practitioners*. 2009;12(2):84-. PMID: 2010218029. Language: English. Entry Date: 20090410. Revision Date: 20090410. Publication Type: journal article. Exclusion Code: X1
17. Anghelescu IG, Kohnen R, Szegedi A, et al. Comparison of Hypericum extract WS 5570 and paroxetine in ongoing treatment after recovery from an episode of moderate to severe depression: results from a randomized multicenter study. *Pharmacopsychiatry*. 2006 Nov;39(6):213-9. PMID: 17124643. Exclusion Code: X2
18. Appelberg BG, Syvälahti EK, Koskinen TE, et al. Patients with severe depression may benefit from buspirone augmentation of selective serotonin reuptake inhibitors: results from a placebo-controlled, randomized, double-blind, placebo wash-in study. *J Clin Psychiatry*; 2001. p. 448-52. Exclusion Code: X5
19. Appelhof BC, Brouwer JP, van Dyck R, et al. Triiodothyronine addition to paroxetine in the treatment of major depressive disorder. *J Clin Endocrinol Metab*. 2004 Dec;89(12):6271-6. PMID: 15579788. Exclusion Code: X2
20. Arnow BA, Blasey C, Manber R, et al. Dropouts versus completers among chronically depressed outpatients. *J Affect Disord*. 2007 Jan;97(1-3):197-202. PMID: 16857266. Exclusion Code: X4
21. Arnow BA, Steidtmann D, Blasey C, et al. The relationship between the therapeutic alliance and treatment outcome in two distinct psychotherapies for chronic depression. *J Consult Clin Psychol*. 2013;81(4):627-38. PMID: 2013-01524-001. PMID: 23339536. First Author & Affiliation: Arnow, Bruce A. Exclusion Code: X2
22. Ashouri A, Atef-Vahid MK, Gharaei B, et al. Effectiveness of meta-cognitive and cognitive-behavioral therapy in patients with major depressive disorder. *Iranian Journal of Psychiatry and Behavioral Sciences*. 2013;7(2):24-34. Exclusion Code: X3
23. Babyak M, Blumenthal JA, Herman S, et al. Exercise treatment for major depression: Maintenance of therapeutic benefit. In: Monat A, Lazarus RS, Reevy G, eds. *The Praeger handbook on stress and coping*. Vol. 2. Westport, CT: Praeger Publishers/Greenwood Publishing Group; 2007:529-40. Exclusion Code: X6
24. Bagby RM, Quilty LC, Segal ZV, et al. Personality and differential treatment response in major depression: a randomized controlled trial comparing cognitive-behavioural therapy and pharmacotherapy. *Can J Psychiatry*. 2008 Jun;53(6):361-70. PMID: 18616856. Exclusion Code: X4
25. Baldomero EB, Ubago JG, Cercós CL, et al. Venlafaxine extended release versus conventional antidepressants in the remission of depressive disorders after previous antidepressant failure: ARGOS study. *Depress Anxiety*. 2005;22(2):68-76. PMID: 2005-15269-004. PMID: 16094658. First Author & Affiliation: Baldomero, E. Baca. Exclusion Code: X3
26. Ballesteros J, Callado LF, Gutierrez M. An independent meta-analysis using summary data for clinical response, remission, and discontinuation for any reason from the 6 pivotal phase III randomized clinical trials of duloxetine in major depressive disorder. *J Clin Psychopharmacol*. 2007 Apr;27(2):219-21. PMID: 17414254. Exclusion Code: X5
27. Bauer M, Bschor T, Kunz D, et al. Double-blind, placebo-controlled trial of the use of lithium to augment antidepressant medication in continuation treatment of unipolar major depression. *Am J Psychiatry*. 2000;157(9):1429-35. Exclusion Code: X2
28. Bauer M, Pretorius HW, Constant EL, et al. Extended-release quetiapine as adjunct to an antidepressant in patients with major depressive disorder: results of a randomized, placebo-controlled, double-blind study. *J Clin Psychiatry*. 2009 Apr;70(4):540-9. PMID: 19358791. Exclusion Code: X3

29. Baumann P, Nil R, Souche A, et al. A double-blind, placebo-controlled study of citalopram with and without lithium in the treatment of therapy-resistant depressive patients: A clinical, pharmacokinetic, and pharmacogenetic investigation. *J Clin Psychopharmacol.* 1996;16(4):307-14. PMID: 1996-05541-004. PMID: 8835706. Partial author list. First Author & Affiliation: Baumann, Pierre. Exclusion Code: X5
30. Bell AC, D'Zurilla TJ. Problem-solving therapy for depression: a meta-analysis. *Clin Psychol Rev.* 2009 Jun;29(4):348-53. PMID: 19299058. Exclusion Code: X2
31. Benazzi F. Fluoxetine and olanzapine for resistant depression. *Am J Psychiatry.* 2002 Jan;159(1):155-6. PMID: 11772722. Exclusion Code: X1
32. Benvenuti A, Rucci P, Miniati M, et al. Treatment-emergent mania/hypomania in unipolar patients. *Bipolar Disord.* 2008 Sep;10(6):726-32. PMID: 18837867. Exclusion Code: X2
33. Berman RM, Darnell AM, Miller HL, et al. Effect of pindolol in hastening response to fluoxetine in the treatment of major depression: a double-blind, placebo-controlled trial. *Am J Psychiatry.* 1997 Jan;154(1):37-43. PMID: 8988956. Exclusion Code: X5
34. Berman RM, Fava M, Thase ME, et al. Aripiprazole augmentation in major depressive disorder: a double-blind, placebo-controlled study in patients with inadequate response to antidepressants. *CNS Spectr.* 2009 Apr;14(4):197-206. PMID: 19407731. Exclusion Code: X5
35. Berman RM, Marcus RN, Swanink R, et al. The efficacy and safety of aripiprazole as adjunctive therapy in major depressive disorder: a multicenter, randomized, double-blind, placebo-controlled study. *J Clin Psychiatry.* 2007 Jun;68(6):843-53. PMID: 17592907. Exclusion Code: X5
36. Bhar SS, Gelfand LA, Schmid SP, et al. Sequence of improvement in depressive symptoms across cognitive therapy and pharmacotherapy. *J Affect Disord.* 2008 Sep;110(1-2):161-6. PMID: 18276017. Exclusion Code: X5
37. Bhattacharya R, Kelley G, Bhattacharjee S. Long-term follow-up effects of computerized or internet-based cognitive behavioral therapy for depression and anxiety: A metaanalysis. *Value Health.* 2012;15(4):A82. Exclusion Code: X5
38. Bisslerbe JC, Boyer P, Souetre E, et al. A Six Month Sertraline Fluoxetine Comparative Study in Depressed Outpatients: Outcome and Costs CONFERENCE ABSTRACT. XXth Collegium Internationale Neuro-psychopharmacologicum. Melbourne, Australia. 23rd-27th June, 1996.; 1996. Exclusion Code: X9
39. Blackburn IM, Bishop S, Glen AI, et al. The efficacy of cognitive therapy in depression: a treatment trial using cognitive therapy and pharmacotherapy, each alone and in combination. *Br J Psychiatry.* 1981 Sep;139:181-9. PMID: 7317698. Exclusion Code: X5
40. Blackburn IM, Moore RG. Controlled acute and follow-up trial of cognitive therapy and pharmacotherapy in out-patients with recurrent depression. *Br J Psychiatry.* 1997 Oct;171:328-34. PMID: 9373420. Exclusion Code: X5
41. Blitzer LE, Atchison-Nevel DJ, Kenny MC. Using acupuncture to treat major depressive disorder: a pilot investigation. *Clinical Acupuncture & Oriental Medicine.* 2003;4(4):144-7. PMID: 2004075709. Language: English. Entry Date: 20040514. Revision Date: 20091218. Publication Type: journal article. Exclusion Code: X4
42. Bloch MH, Hannestad J. Omega-3 fatty acids for the treatment of depression: systematic review and meta-analysis. *Mol Psychiatry.* 2012 Dec;17(12):1272-82. PMID: 21931319. Exclusion Code: X5
43. Blom MB, Hoek HW, Spinhoven P, et al. Treatment of depression in patients from ethnic minority groups in the Netherlands. *Transcult Psychiatry.* 2010 Jul;47(3):473-90. PMID: 20688800. Exclusion Code: X4
44. Blom MBJ, Spinhoven P, Hoffman T, et al. Severity and duration of depression, not personality factors, predict short term outcome in the treatment of major depression. *J Affect Disord.* 2007;104(1-3):119-26. Exclusion Code: X6

45. Bobo WV, Chen H, Trivedi MH, et al. Randomized comparison of selective serotonin reuptake inhibitor (escitalopram) monotherapy and antidepressant combination pharmacotherapy for major depressive disorder with melancholic features: a CO-MED report. *J Affect Disord*. 2011 Oct;133(3):467-76. PMID: 21601287. Exclusion Code: X5
46. Bobo WV, Shelton RC. Fluoxetine and olanzapine combination therapy in treatment-resistant major depression: review of efficacy and safety data. *Expert Opin Pharmacother*. 2009 Sep;10(13):2145-59. PMID: 19640209. Exclusion Code: X1
47. Bolton PG, Fergusson KJ, Parker SM, et al. Randomised controlled trial of cognitive-behavioural therapy and routine GP care for major depression. *Med J Aust*. 2001 Jul 16;175(2):118-9. PMID: 11556412. Exclusion Code: X5
48. Bone K. St. John's Wort and depression: Avoiding drug interactions. *Townsend Letter for Doctors and Patients*. 2007;282:45-8. PMID: 0094709. Exclusion Code: X1
49. Bortolotti B, Menchetti M, Bellini F, et al. Psychological interventions for major depression in primary care: a meta-analytic review of randomized controlled trials. *Gen Hosp Psychiatry*. 2008 Jul-Aug;30(4):293-302. PMID: 18585531. Exclusion Code: X5
50. Bosmans JE, van Schaik DJ, Heymans MW, et al. Cost-effectiveness of interpersonal psychotherapy for elderly primary care patients with major depression. *Int J Technol Assess Health Care*. 2007 Fall;23(4):480-7. PMID: 17937837. Exclusion Code: X5
51. Bosscher RJ. Running and mixed physical exercises with depressed psychiatric patients. *Int J Sport Psychol*. 1993;24(2):170-84. PMID: 1994-06914-001. First Author & Affiliation: Bosscher, Rudolf J. Exclusion Code: X2
52. Braun SR, Gregor B, Tran US. Comparing bona fide psychotherapies of depression in adults with two meta-analytical approaches. *PLoS One*. 2013;8(6)PMID: 2013-25847-001. First Author & Affiliation: Braun, Sarah R. Exclusion Code: X5
53. Brenner R, Madhusoodanan S, Pawlowska M. Efficacy of continuation treatment with *Hypericum perforatum* in depression. *J Clin Psychiatry*. 2002;63(5)PMID: 2002-13479-018. PMID: 12019675. First Author & Affiliation: Brenner, Ronald. Exclusion Code: X4
54. Brody A, Saxena S, Stoessel P, et al. Regional brain metabolic changes in patients with major depression treated with either paroxetine or interpersonal therapy. *Arch Gen Psychiatry*. 2001 Jul;58(7):631-40. PMID: 0028127. Exclusion Code: X4
55. Brosse A, Sheets E, Lett H, et al. Exercise and the treatment of clinical depression in adults: recent findings and future directions. *Sports Medicine (Auckland)*. 2002;32(12):741-60. PMID: 0043625. Exclusion Code: X4
56. Brown WA, Harrison W. Are patients who are intolerant to one SSRI intolerant to another? *Psychopharmacol Bull*. 1992;28(3):253-6. PMID: 1480728. Exclusion Code: X5
57. Brown WA, Harrison W. Are patients who are intolerant to one serotonin selective reuptake inhibitor intolerant to another? *J Clin Psychiatry*. 1995 Jan;56(1):30-4. PMID: 7836337. Exclusion Code: X5
58. Bulmash E, Harkness KL, Stewart JG, et al. Personality, stressful life events, and treatment response in major depression. *J Consult Clin Psychol*. 2009 Dec;77(6):1067-77. PMID: 19968383. Exclusion Code: X5
59. Buysse DJ, Kupfer DJ, Cherry C, et al. Effects of prior fluoxetine treatment on EEG sleep in women with recurrent depression. *Neuropsychopharmacology*. 1999 Aug;21(2):258-67. PMID: 10432474. Exclusion Code: X4
60. Carpenter LL, Yasmin S, Price LH. A double-blind, placebo-controlled study of mirtazapine augmentation for refractory major depression: preliminary findings. *52nd Institute on Psychiatric Services*; 2000 October 25-29th; Philadelphia, PA; 2000. Exclusion Code: X9

61. Carpenter LL, Yasmin S, Price LH. A double-blind, placebo-controlled study of antidepressant augmentation with mirtazapine. *Biol Psychiatry*. 2002 Jan 15;51(2):183-8. PMID: 11822997. Exclusion Code: X5
62. Carta MG, Hardoy MC, Pulu A, et al. Improving physical quality of life with group physical activity in the adjunctive treatment of major depressive disorder. *Clin Pract Epidemiol Ment Health*. 2008;4PMID: 2008-15806-001. First Author & Affiliation: Carta, Mauro Giovanni. Exclusion Code: X5
63. Carvalho AF, Berk M, Hyphantis TN, et al. The integrative management of treatment-resistant depression: a comprehensive review and perspectives. *Psychother Psychosom*. 2014;83(2):70-88. PMID: 24458008. Exclusion Code: X2
64. Casacalenda N, Perry JC, Looper K. Remission in major depressive disorder: a comparison of pharmacotherapy, psychotherapy, and control conditions. *Am J Psychiatry*. 2002 Aug;159(8):1354-60. PMID: 12153828. Exclusion Code: X3
65. Chalder M, Wiles NJ, Campbell J, et al. Facilitated physical activity as a treatment for depressed adults: Randomised controlled trial. *BMJ: British Medical Journal*. 2012;344(7860)PMID: 2012-15758-001. First Author & Affiliation: Chalder, Melanie. Exclusion Code: X5
66. Chen J, Gao K, Kemp DE. Second-generation antipsychotics in major depressive disorder: update and clinical perspective. *Curr Opin Psychiatry*. 2011 Jan;24(1):10-7. PMID: 21088586. Exclusion Code: X3
67. Chiesa A, Serretti A. L'utilità delle meditazioni basate sulla consapevolezza per i disturbi psichiatrici: Una review sistematica. *Psichiatria e Psicoterapia*. 2009;28(2):93-110. PMID: 2009-17997-001. First Author & Affiliation: Chiesa, Alberto. Translated Title: Usefulness of mindfulness meditations for psychiatric disorders: A systematic review.. Other Journal Titles: *Psichiatria e Psicoterapia Analitica*. Release Date: 20110110. Publication Type: Journal, (0100). Exclusion Code: X1
68. Chiesa A, Serretti A. Mindfulness based cognitive therapy for psychiatric disorders: a systematic review and meta-analysis. *Psychiatry Res*. 2011 May 30;187(3):441-53. PMID: 20846726. Exclusion Code: X3
69. Cody RA, Drysdale K. The effects of psychotherapy on reducing depression in residential aged care: A meta-analytic review. *Clinical Gerontologist: The Journal of Aging and Mental Health*. 2013;36(1):46-69. PMID: 2012-35079-004. First Author & Affiliation: Cody, Robyn Ann. Exclusion Code: X5
70. Coelho HF, Canter PH, Ernst E. Mindfulness-based cognitive therapy: evaluating current evidence and informing future research. *J Consult Clin Psychol*. 2007 Dec;75(6):1000-5. PMID: 18085916. Exclusion Code: X5
71. Combs K, Smith PJ, Sherwood A, et al. Impact of sleep complaints and depression outcomes among participants in the standard medical intervention and long-term exercise study of exercise and pharmacotherapy for depression. *J Nerv Ment Dis*. 2014 Feb;202(2):167-71. PMID: 24469530. Exclusion Code: X4
72. Coon DW, DeVries HM, Gallagher-Thompson D. COGNITIVE BEHAVIORAL THERAPY WITH SUICIDAL OLDER ADULTS. *Behav Cogn Psychother*. 2004;32(04):481-93. Exclusion Code: X1
73. Cooney GM, Dwan K, Greig CA, et al. Exercise for depression. *Cochrane Database Syst Rev*. 2013;9:CD004366. PMID: 24026850. Exclusion Code: X5
74. Cornwall PL, Jenaway A, Garland A, et al. Cognitive therapy for major depression in partial remission: preliminary findings. 9th Congress of the Association of European Psychiatrists. Copenhagen, Denmark. 20-24th September 1998.; 1998. Exclusion Code: X9
75. Corya SA, Williamson D, Sanger TM, et al. A randomized, double-blind comparison of olanzapine/fluoxetine combination, olanzapine, fluoxetine, and venlafaxine in treatment-resistant depression. *Depress Anxiety*. 2006;23(6):364-72. PMID: 16710853. Exclusion Code: X2

76. Cuijpers P, de Beurs DP, van Spijker BAJ, et al. The effects of psychotherapy for adult depression on suicidality and hopelessness: A systematic review and meta-analysis. *J Affect Disord.* 2013;144(3):183-90. PMID: 2012-20095-001. PMID: 22832172. First Author & Affiliation: Cuijpers, Pim. Exclusion Code: X6
77. Cuijpers P, Dekker J, Hollon SD, et al. Adding psychotherapy to pharmacotherapy in the treatment of depressive disorders in adults: a meta-analysis. *J Clin Psychiatry.* 2009 Sep;70(9):1219-29. PMID: 19818243. Exclusion Code: X3
78. Cuijpers P, Reynolds CF, III, Donker T, et al. Personalized treatment of adult depression: Medication, psychotherapy, or both? A systematic review. *Depress Anxiety.* 2012;29(10):855-64. PMID: 2012-26844-004. PMID: 22815247. First Author & Affiliation: Cuijpers, Pim. Exclusion Code: X2
79. Cuijpers P, Sijbrandij M, Koole SL, et al. Adding psychotherapy to antidepressant medication in depression and anxiety disorders: a meta-analysis. *World Psychiatry.* 2014 Feb;13(1):56-67. PMID: 24497254. Exclusion Code: X5
80. Cuijpers P, Straten A, Schuurmans J, et al. Psychotherapy for chronic major depression and dysthymia: a meta-analysis (Structured abstract). *Clin Psychol Rev;* 2009. p. 51-62. Exclusion Code: X5
81. Cuijpers P, Straten A, Warmerdam L. Behavioral activation treatments of depression: a meta-analysis (Structured abstract). *Clin Psychol Rev;* 2007. p. 318-26. Exclusion Code: X2
82. Cuijpers P, van Straten A, Hollon SD, et al. The contribution of active medication to combined treatments of psychotherapy and pharmacotherapy for adult depression: A meta-analysis. *Acta Psychiatr Scand.* 2010;121(6):415-23. PMID: 2010-09276-003. PMID: 19922522. First Author & Affiliation: Cuijpers, P. Exclusion Code: X2
83. Cuijpers P, van Straten A, Smit F, et al. Is psychotherapy for depression equally effective in younger and older adults? A meta-regression analysis. *Int Psychogeriatr.* 2009;21(1):16-24. PMID: 2009-02458-005. PMID: 19040783. First Author & Affiliation: Cuijpers, Pim. Exclusion Code: X5
84. Cuijpers P, van Straten A, van Oppen P, et al. Are psychological and pharmacologic interventions equally effective in the treatment of adult depressive disorders? A meta-analysis of comparative studies. *J Clin Psychiatry.* 2008 Nov;69(11):1675-85; quiz 839-41. PMID: 18945396. Exclusion Code: X5
85. Cuijpers P, van Straten A, van Oppen P, et al. Comparing psychotherapy and pharmacotherapy for adult depression: adjusting for differential dropout rates. *J Clin Psychiatry.* 2010 Sep;71(9):1246. PMID: 20923628. Exclusion Code: X1
86. Cuijpers P, van Straten A, Warmerdam L, et al. Psychological treatment of depression: A meta-analytic database of randomized studies. *BMC Psychiatry.* 2008;8PMID: 2008-10088-001. PMID: 18485191. First Author & Affiliation: Cuijpers, Pim. Exclusion Code: X1
87. Cuijpers P, van Straten A, Warmerdam L, et al. Psychotherapy versus the combination of psychotherapy and pharmacotherapy in the treatment of depression: a meta-analysis. *Depress Anxiety.* 2009;26(3):279-88. PMID: 19031487. Exclusion Code: X3
88. Curran S, Rampat R, Spencer R, et al. Paroxetine for the treatment of old age depression: A systematic review. *Eur Neuropsychopharmacol.* 2009;19:S403-S4. Exclusion Code: X1
89. Daley A. Exercise and depression: A review of reviews. *J Clin Psychol Med Settings.* 2008;15(2):140-7. PMID: 19104978. Exclusion Code: X2
90. Dannlowski U, Baune BT, Bockermann I, et al. Adjunctive antidepressant treatment with quetiapine in agitated depression: Positive effects on symptom reduction, psychopathology and remission rates. *Human Psychopharmacology.* 2008;23(7):587-93. Exclusion Code: X4

91. Davis AS. A meta-analysis of the efficacy of cognitive therapy, pharmacotherapy, and the combination of cognitive therapy and pharmacotherapy in the treatment of depression. US: ProQuest Information & Learning; 1991. Exclusion Code: X9
92. de Graaf LE, Gerhards SAH, Arntz A, et al. One-year follow-up results of unsupported online computerized cognitive behavioural therapy for depression in primary care: A randomized trial. *J Behav Ther Exp Psychiatry*. 2011;42(1):89-95. PMID: 2010-26707-013. PMID: 20723885. First Author & Affiliation: de Graaf, L. E. Exclusion Code: X2
93. de Jonghe F, Hendricksen M, van Aalst G, et al. Psychotherapy alone and combined with pharmacotherapy in the treatment of depression. *Br J Psychiatry*. 2004 Jul;185:37-45. PMID: 15231554. Exclusion Code: X5
94. de Jonghe F, Kool S, van Aalst G, et al. Combining psychotherapy and antidepressants in the treatment of depression. *J Affect Disord*. 2001 May;64(2-3):217-29. PMID: 11313088. Exclusion Code: X5
95. De Jong-Meyer R, Hautzinger M, Rudolf GAE, et al. Outcome prediction and longitudinal analyses of endogenously depressed patients treated with combined psychological and antidepressant therapies. *Zeitschrift fur Klinische Psychologie*. 1996;25(2):110-29. Exclusion Code: X1
96. de la Cerda P, Cervello E, Cocca A, et al. Effect of an aerobic training program as complementary therapy in patients with moderate depression. *Percept Mot Skills*. 2011 Jun;112(3):761-9. PMID: 21853765. Exclusion Code: X4
97. De Maat S, Dekker J, Schoevers R, et al. Relative efficacy of psychotherapy and pharmacotherapy in the treatment of depression: A meta-analysis. *Psychotherapy Research*. 2006;16(5):562-72. PMID: 2006-21953-005. First Author & Affiliation: De Maat, Saskia. Exclusion Code: X5
98. de Maat S, Dekker J, Schoevers R, et al. Short psychodynamic supportive psychotherapy, antidepressants, and their combination in the treatment of major depression: a mega-analysis based on three randomized clinical trials. *Depress Anxiety*. 2008;25(7):565-74. PMID: 17557313. Exclusion Code: X4
99. de Mello MF, de Jesus Mari J, Bacaltchuk J, et al. A systematic review of research findings on the efficacy of interpersonal therapy for depressive disorders. *Eur Arch Psychiatry Clin Neurosci*. 2005 Apr;255(2):75-82. PMID: 15812600. Exclusion Code: X5
100. DeBattista C, Doghramji K, Menza MA, et al. Adjunct modafinil for the short-term treatment of fatigue and sleepiness in patients with major depressive disorder: A preliminary double-blind, placebo-controlled study. *J Clin Psychiatry*. 2003;64(9):1057-64. Exclusion Code: X5
101. DeBattista C, Solvason HB, Poirier J, et al. A prospective trial of bupropion SR augmentation of partial and non-responders to serotonergic antidepressants. *J Clin Psychopharmacol*. 2003 Feb;23(1):27-30. PMID: 12544372. Exclusion Code: X5
102. Den Boer PCAM, Wiersma D, Ten Vaarwerk I, et al. Cognitive self-therapy for chronic depression and anxiety: A multi-centre randomized controlled study. *Psychological Medicine: A Journal of Research in Psychiatry and the Allied Sciences*. 2007;37(3):329-39. PMID: 2007-03729-003. First Author & Affiliation: Den Boer, Peter C. A. M. Exclusion Code: X2
103. Denton WH, Carmody TJ, Rush AJ, et al. Dyadic discord at baseline is associated with lack of remission in the acute treatment of chronic depression. *Psychol Med*. 2010 Mar;40(3):415-24. PMID: 19607755. Exclusion Code: X2
104. DeRubeis RJ, Evans MD, Hollon SD, et al. How does cognitive therapy work? Cognitive change and symptom change in cognitive therapy and pharmacotherapy for depression. *J Consult Clin Psychol*. 1990 Dec;58(6):862-9. PMID: 2292637. Exclusion Code: X5

105. DeRubeis RJ, Gelfand LA, Tang TZ, et al. Medications versus cognitive behavior therapy for severely depressed outpatients: Mega-analysis of four randomized comparisons. *The American Journal of Psychiatry*. 1999;156(7):1007-13. PMID: 1999-03084-003. PMID: 10401443. First Author & Affiliation: DeRubeis, Robert J. Exclusion Code: X4
106. Dew MA, Whyte EM, Lenze EJ, et al. Recovery from major depression in older adults receiving augmentation of antidepressant pharmacotherapy. *Am J Psychiatry*. 2007;164(6):892-9. PMID: 2009673818. Language: English. Entry Date: 20071012. Revision Date: 20091218. Publication Type: journal article. Exclusion Code: X5
107. Dhillon S, Yang LPH, Curran MP. Spotlight on bupropion in major depressive disorder. *CNS Drugs*. 2008;22(7):613-7. PMID: 2008-08817-006. PMID: 18547129. First Author & Affiliation: Dhillon, Sohita. Exclusion Code: X1
108. Dombrovski AY, Lenze EJ, Dew MA, et al. Maintenance treatment for old-age depression preserves health-related quality of life: a randomized, controlled trial of paroxetine and interpersonal psychotherapy. *J Am Geriatr Soc*. 2007 Sep;55(9):1325-32. PMID: 17767673. Exclusion Code: X2
109. Dording CM, Mischoulon D, Shyu I, et al. SAMe and sexual functioning. *Eur Psychiatry*. 2012 Aug;27(6):451-4. PMID: 21398094. Exclusion Code: X3
110. Doree JP, Des Rosiers J, Lew V, et al. Quetiapine augmentation of treatment-resistant depression: A comparison with lithium. *Curr Med Res Opin*. 2007;23(2):333-41. Exclusion Code: X2
111. Dozois DJ, Bieling PJ, Patelis-Siotis I, et al. Changes in self-schema structure in cognitive therapy for major depressive disorder: a randomized clinical trial. *J Consult Clin Psychol*. 2009 Dec;77(6):1078-88. PMID: 19968384. Exclusion Code: X2
112. Dunlop BW, Crits-Christoph P, Evans DL, et al. Coadministration of modafinil and a selective serotonin reuptake inhibitor from the initiation of treatment of major depressive disorder with fatigue and sleepiness: A double-blind, placebo-controlled study. *J Clin Psychopharmacol*. 2007;27(6):614-9. Exclusion Code: X5
113. Dunlop BW, Kelley ME, Mletzko TC, et al. Depression beliefs, treatment preference, and outcomes in a randomized trial for major depressive disorder. *J Psychiatr Res*. 2012 Mar;46(3):375-81. PMID: 22118808. Exclusion Code: X7
114. Dunner DL, Amsterdam JD, Shelton RC, et al. Efficacy and tolerability of adjunctive ziprasidone in treatment-resistant depression: a randomized, open-label, pilot study. *J Clin Psychiatry*. 2007 Jul;68(7):1071-7. PMID: 17685744. Exclusion Code: X2
115. Eisendrath SJ, Gillung E, Delucchi K, et al. A preliminary study: Efficacy of mindfulness-based cognitive therapy versus sertraline as first-line treatments for major depressive disorder. *Mindfulness*. 2014PMID: 2014-03529-001. Publication Status: Online First Posting. First Author & Affiliation: Eisendrath, Stuart J.. Release Date: 20140203. Publication Type: Journal, (0100). Exclusion Code: X4
116. Ekers D, Richards D, Gilbody S. A meta-analysis of randomized trials of behavioural treatment of depression. *Psychological Medicine: A Journal of Research in Psychiatry and the Allied Sciences*. 2008;38(5):611-23. PMID: 2008-05402-002. First Author & Affiliation: Ekers, D. Exclusion Code: X5
117. Ekselius L, von Knorring L. Effect on sexual function of long-term treatment with selective serotonin reuptake inhibitors in depressed patients treated in primary care. *J Clin Psychopharmacol*. 2001 Apr;21(2):154-60. PMID: 11270911. Exclusion Code: X3

118. El-Khalili N, Joyce M, Atkinson S, et al. Extended-release quetiapine fumarate (quetiapine XR) as adjunctive therapy in major depressive disorder (MDD) in patients with an inadequate response to ongoing antidepressant treatment: A multicentre, randomized, double-blind, placebo-controlled study. *Int J Neuropsychopharmacol*. 2010;13(7):917-32. Exclusion Code: X5
119. Eriksson E. Cognitive behavioural therapy for treatment-resistant depression. *Lancet*. 2013 May 25;381(9880):1814-5. PMID: 0169677. Exclusion Code: X1
120. Eriksson S, Gard G. Physical exercise and depression. *Phys Ther Rev*. 2011 Aug;16(4):261-8. PMID: 0148106. Exclusion Code: X4
121. Ernst E, Lee MS, Choi T-Y. Acupuncture for depression?: A systematic review of systematic reviews. *Eval Health Prof*. 2011;34(4):403-12. PMID: 2011-29673-001. First Author & Affiliation: Ernst, E. Exclusion Code: X4
122. Fang Y, Yuan C, Xu Y, et al. A pilot study of the efficacy and safety of paroxetine augmented with risperidone, valproate, buspirone, trazodone, or thyroid hormone in adult Chinese patients with treatment-resistant major depression. *J Clin Psychopharmacol*. 2011 Oct;31(5):638-42. PMID: 21869688. Exclusion Code: X2
123. Farabaugh A, Alpert J, Wisniewski SR, et al. Cognitive therapy for anxious depression in STAR(*) D: what have we learned? *J Affect Disord*. 2012 Dec 15;142(1-3):213-8. PMID: 22877961. Exclusion Code: X4
124. Fava GA, Rafanelli C, Grandi S, et al. Prevention of recurrent depression with cognitive behavioral therapy: preliminary findings. *Arch Gen Psychiatry*. 1998 Sep;55(9):816-20. PMID: 9736008. Exclusion Code: X2
125. Fava GA, Ruini C, Rafanelli C, et al. Cognitive behavior approach to loss of clinical effect during long-term antidepressant treatment: a pilot study. *Am J Psychiatry*. 2002 Dec;159(12):2094-5. PMID: 12450962. Exclusion Code: X2
126. Fava M, Alpert J, Nierenberg A, et al. Double-blind study of high-dose fluoxetine versus lithium or desipramine augmentation of fluoxetine in partial responders and nonresponders to fluoxetine. *J Clin Psychopharmacol*. 2002 Aug;22(4):379-87. PMID: 12172337. Exclusion Code: X5
127. Fava M, Rosenbaum JF, McGrath PJ, et al. Lithium and tricyclic augmentation of fluoxetine treatment for resistant major depression: a double-blind, controlled study. *Am J Psychiatry*. 1994 Sep;151(9):1372-4. PMID: 8067495. Exclusion Code: X5
128. Fava M, Rush AJ, Alpert JE, et al. Difference in treatment outcome in outpatients with anxious versus nonanxious depression: a STAR*D report. *Am J Psychiatry*. 2008 Mar;165(3):342-51. PMID: 18172020. Exclusion Code: X2
129. Feldman G, Harley R, Kerrigan M, et al. Change in emotional processing during a dialectical behavior therapy-based skills group for major depressive disorder. *Behav Res Ther*. 2009 Apr;47(4):316-21. PMID: 19232571. Exclusion Code: X5
130. Feng CY, Chu H, Chen CH, et al. The effect of cognitive behavioral group therapy for depression: A meta-analysis 2000–2010. *Worldviews Evid Based Nurs*. 2012;9(1):2-17. PMID: 2012-03102-002. PMID: 22221447. First Author & Affiliation: Feng, Chiueng-Yi. Exclusion Code: X5
131. Finch EJJ, Katona CLE. Lithium augmentation in the treatment of refractory depression in old age. *Int J Geriatr Psychiatry*. 1989;4(1):41-6. Exclusion Code: X3
132. Fjorback LO, Arendt M, Ørnbøl E, et al. Mindfulness-Based Stress Reduction and Mindfulness-Based Cognitive Therapy—A systematic review of randomized controlled trials. *Acta Psychiatr Scand*. 2011;124(2):102-19. PMID: 2011-14452-003. PMID: 21534932. First Author & Affiliation: Fjorback, L. O. Exclusion Code: X2
133. Fleurence R, Williamson R, Jing Y, et al. A systematic review of augmentation strategies for patients with major depressive disorder. *Psychopharmacol Bull*. 2009;42(3):57-90. PMID: 19752841. Exclusion Code: X3

134. Flint AJ, Rifat SL. The effect of sequential antidepressant treatment on geriatric depression. *J Affect Disord.* 1996 Jan 22;36(3-4):95-105. PMID: 8821312. Exclusion Code: X5
135. Forest L. Two-Week Double-Blind Placebo-Controlled Study of Escitalopram in the Treatment of Severe Major Depression [SCT-MD-26]. Forest Laboratories - Clinical Study Register [www.forestclinicaltrials.com]; 2002. Exclusion Code: X9
136. Fournier JC, DeRubeis RJ, Hollon SD, et al. Antidepressant drug effects and depression severity: a patient-level meta-analysis. *JAMA.* 2010 Jan 6;303(1):47-53. PMID: 20051569. Exclusion Code: X5
137. Fournier JC, DeRubeis RJ, Hollon SD, et al. Differential change in specific depressive symptoms during antidepressant medication or cognitive therapy. *Behav Res Ther.* 2013;51(7):392-8. PMID: 2013-19059-010. PMID: 23644038. First Author & Affiliation: Fournier, Jay C. Exclusion Code: X4
138. Fournier JC, DeRubeis RJ, Shelton RC, et al. Prediction of response to medication and cognitive therapy in the treatment of moderate to severe depression. *J Consult Clin Psychol.* 2009 Aug;77(4):775-87. PMID: 19634969. Exclusion Code: X7
139. Freeman MP, Hibbeln JR, Wisner KL, et al. Omega-3 fatty acids: evidence basis for treatment and future research in psychiatry. *J Clin Psychiatry.* 2006 Dec;67(12):1954-67. PMID: 17194275. Exclusion Code: X1
140. Freeman MP, Mischoulon D, Tedeschini E, et al. Complementary and alternative medicine for major depressive disorder: a meta-analysis of patient characteristics, placebo-response rates, and treatment outcomes relative to standard antidepressants (Provisional abstract). *J Clin Psychiatry*; 2010. p. 682-8. Exclusion Code: X5
141. Freeman MP, Rapaport MH. Omega-3 fatty acids and depression: from cellular mechanisms to clinical care. *J Clin Psychiatry.* 2011 Feb;72(2):258-9. PMID: 21382308. Exclusion Code: X1
142. Friede M, Henneicke von Zepelin HH, Freudenstein J. Differential therapy of mild to moderate depressive episodes (ICD-10 F 32.0; F 32.1) with St. John's wort. *Pharmacopsychiatry.* 2001;34(Suppl1):S38-S41. PMID: 2001-11122-006. First Author & Affiliation: Friede, Michael. Exclusion Code: X4
143. Gahlsdorf T, Krause R, Beal M. Efficacy of St John's Wort for treating mild to moderate depression. *Complementary Health Practice Review.* 2007 Oct;12(3):184-95. PMID: 0105158. Exclusion Code: X8
144. Gangadhar BN, Naveen GH, Rao MG, et al. Positive antidepressant effects of generic yoga in depressive out-patients: A comparative study. *Indian J Psychiatry.* 2013;55(7):S369-S73. Exclusion Code: X3
145. Garakani A, Martinez JM, Marcus S, et al. A randomized, double-blind, and placebo-controlled trial of quetiapine augmentation of fluoxetine in major depressive disorder. *Int Clin Psychopharmacol.* 2008 Sep;23(5):269-75. PMID: 18703936. Exclusion Code: X5
146. Gartlehner G, Thieda P, Hansen RA, et al. Comparative risk for harms of second-generation antidepressants : a systematic review and meta-analysis. *Drug Saf.* 2008;31(10):851-65. PMID: 18759509. Exclusion Code: X2
147. Gilbert G. Adults with both anxiety and depression respond poorly to treatment. *J Natl Med Assoc.* 2008;100(7):870-1. Exclusion Code: X1
148. Gloaguen V, Cottraux J, Cucherat M, et al. A meta-analysis of the effects of cognitive therapy in depressed patients. *J Affect Disord.* 1998 Apr;49(1):59-72. PMID: 9574861. Exclusion Code: X5
149. Goldstein CN, Topel ME, Zajecka JM. Second generation antipsychotic medications in the treatment of affective disorders. In: Schwartz TL, Megna J, Topel ME, eds. *Antipsychotic drugs: Pharmacology, side effects and abuse prevention.* Hauppauge, NY US: Nova Biomedical Books; 2013:9-31. Exclusion Code: X1

150. Gould R, Coulson M, Howard R. Cognitive behavioral therapy for depression in older people: A meta-analysis and meta-regression of randomized controlled trials. *Journal - American Geriatrics Society*. 2012 Oct;60(10):1817-30. PMID: 0161072. Exclusion Code: X5
151. Green BL, Krupnick JL, Chung J, et al. Impact of PTSD comorbidity on one-year outcomes in a depression trial. *J Clin Psychol*. 2006 Jul;62(7):815-35. PMID: 16703602. Exclusion Code: X2
152. Greenlee A, Karp JF, Dew MA, et al. Anxiety impairs depression remission in partial responders during extended treatment in late-life. *Depress Anxiety*. 2010 May;27(5):451-6. PMID: 20186975. Exclusion Code: X5
153. Grobler AC, Matthews G, Molenberghs G. The impact of missing data on clinical trials: a re-analysis of a placebo controlled trial of *Hypericum perforatum* (St Johns wort) and sertraline in major depressive disorder. *Psychopharmacology (Berl)*. 2013 Nov 15; PMID: 24232445. Exclusion Code: X4
154. Güemes I, Guillén V, Ballesteros J. Psychotherapy versus drug therapy in depression in outpatient care. *Actas Españolas de Psiquiatría*. 2008;36(5):299-306. PMID: 2008-16648-007. PMID: 18523896. First Author & Affiliation: Güemes, I. Exclusion Code: X1
155. Gulrez G, Badyal DK, Deswal RS, et al. Bupropion as an augmenting agent in patients of depression with partial response. *Basic Clin Pharmacol Toxicol*. 2012 Mar;110(3):227-30. PMID: 21895979. Exclusion Code: X5
156. Haeffel G. CBT self-help that harms. *Human Givens*. 2010;17(2):8. PMID: 0133238. Exclusion Code: X1
157. Han C, Wang SM, Seo HJ, et al. Aripiprazole augmentation, antidepressant combination or switching therapy in patients with major depressive disorder who are partial- or non-responsive to current antidepressants: A multi-center, naturalistic study. *J Psychiatr Res*. 2014;49(1):75-82. Exclusion Code: X5
158. Harkness KL, Bagby RM, Kennedy SH. Childhood maltreatment and differential treatment response and recurrence in adult major depressive disorder. *J Consult Clin Psychol*. 2012 Jun;80(3):342-53. PMID: 22428942. Exclusion Code: X5
159. Hecht H, van Calker D. [Evidence-based psychotherapy of depressive disorders: acute therapy and maintenance therapy]. *Psychother Psychosom Med Psychol*. 2008 Aug;58(8):326-34; quiz 35-8. PMID: 18626847. Exclusion Code: X1
160. Herman S, Blumenthal JA, Babyak M, et al. Exercise therapy for depression in middle-aged and older adults: predictors of early dropout and treatment failure. *Health Psychol*. 2002 Nov;21(6):553-63. PMID: 12433007. Exclusion Code: X4
161. Hermens ML, van Hout HP, Terluin B, et al. Clinical effectiveness of usual care with or without antidepressant medication for primary care patients with minor or mild-major depression: a randomized equivalence trial. *BMC Med*. 2007;5:36. PMID: 18067659. Exclusion Code: X3
162. Hinrichsen GA. Treating older adults with interpersonal psychotherapy for depression. *J Clin Psychol*. 1999;55(8):949-60. Exclusion Code: X1
163. Hirschfeld RM, Dunner DL, Keitner G, et al. Does psychosocial functioning improve independent of depressive symptoms? A comparison of nefazodone, psychotherapy, and their combination. *Biol Psychiatry*. 2002 Jan 15;51(2):123-33. PMID: 11822991. Exclusion Code: X2
164. Hirvonen J, Hietala J, Kajander J, et al. Effects of antidepressant drug treatment and psychotherapy on striatal and thalamic dopamine D2/3 receptors in major depressive disorder studied with [11C]raclopride PET. *J Psychopharmacol*. 2011 Oct;25(10):1329-36. PMID: 20829308. Exclusion Code: X5
165. Hoffman BM, Babyak MA, Craighead WE, et al. Exercise and pharmacotherapy in patients with major depression: one-year follow-up of the SMILE study. *Psychosom Med*. 2011 Feb-Mar;73(2):127-33. PMID: 21148807. Exclusion Code: X4

166. Hofmann SG, Sawyer AT, Witt AA, et al. The effect of mindfulness-based therapy on anxiety and depression: A meta-analytic review. *J Consult Clin Psychol*. 2010 Apr;78(2):169-83. PMID: 20350028. Exclusion Code: X5
167. Høifødt RS, Strøm C, Kolstrup N, et al. Effectiveness of cognitive behavioural therapy in primary health care: A review. *Fam Pract*. 2011;28(5):489-504. PMID: 2011-24876-005. PMID: 21555339. First Author & Affiliation: Høifødt, Ragnhild Sørensen. Exclusion Code: X2
168. Hollon SD, Jarrett RB, Nierenberg AA, et al. Psychotherapy and medication in the treatment of adult and geriatric depression: Which monotherapy or combined treatment? *J Clin Psychiatry*. 2005;66(4):455-68. Exclusion Code: X1
169. Hsiao FH, Jow GM, Lai YM, et al. The long-term effects of psychotherapy added to pharmacotherapy on morning to evening diurnal cortisol patterns in outpatients with major depression. *Psychother Psychosom*. 2011;80(3):166-72. PMID: 21389753. Exclusion Code: X3
170. İlhan Yargic L, Corapcioglu A, Kocabasoglu N, et al. A prospective randomized single-blind, multicenter trial comparing the efficacy and safety of paroxetine with and without quetiapine therapy in depression associated with anxiety. *Int J Psychiatry Clin Pract*. 2004;8(4):205-11. Exclusion Code: X5
171. Jakobsen JC, Hansen JL, Simonsen E, et al. The effect of adding psychodynamic therapy to antidepressants in patients with major depressive disorder. A systematic review of randomized clinical trials with meta-analyses and trial sequential analyses (Provisional abstract). *Database of Abstracts of Reviews of Effects*; 2012. p. 4-14. Exclusion Code: X5
172. Jakobsen JC, Hansen JL, Simonsen S, et al. Effects of cognitive therapy versus interpersonal psychotherapy in patients with major depressive disorder: a systematic review of randomized clinical trials with meta-analyses and trial sequential analyses. *Psychol Med*. 2012 Jul;42(7):1343-57. PMID: 22051174. Exclusion Code: X5
173. Jakobsen JC, Hansen JL, Storebo OJ, et al. The effects of cognitive therapy versus 'treatment as usual' in patients with major depressive disorder. *PLoS One*; 2011. Exclusion Code: X5
174. Jarrett RB, Minhajuddin A, Gershenfeld H, et al. Preventing depressive relapse and recurrence in higher-risk cognitive therapy responders: a randomized trial of continuation phase cognitive therapy, fluoxetine, or matched pill placebo. *JAMA Psychiatry*. 2013 Nov;70(11):1152-60. PMID: 24005123. Exclusion Code: X2
175. Jarrett RB, Thase ME. Comparative efficacy and durability of continuation phase cognitive therapy for preventing recurrent depression: design of a double-blinded, fluoxetine- and pill placebo-controlled, randomized trial with 2-year follow-up. *Contemp Clin Trials*. 2010 Jul;31(4):355-77. PMID: 20451668. Exclusion Code: X2
176. Joyce AS, O'Kelly JG, Ogrodniczuk JS, et al. A naturalistic trial of brief psychodynamic therapy for recurrent major depression. *Psychodyn Psychiatry*. 2012 Dec;40(4):645-71. PMID: 23216401. Exclusion Code: X5
177. Kaltenthaler E, Parry G, Beverley C. Computerized Cognitive Behaviour Therapy: A Systematic Review. *Behav Cogn Psychother*. 2004;32(1):31-55. PMID: 2004-12191-003. First Author & Affiliation: Kaltenthaler, Eva. Exclusion Code: X2
178. Kamath J. A Randomized, Double-Blind Placebo-controlled Study Evaluating the Efficacy of Omega 3 Fatty Acid Augmentation of Desvenlafaxine for the Treatment of Major Depressive Disorder in Patients With Medical Illness. [Http://clinicaltrials.gov/show/NCT01803711](http://clinicaltrials.gov/show/NCT01803711) ; 2013. Exclusion Code: X9
179. Karlsson H, Hirvonen J, Salminen J, et al. Increased serotonin receptor 1A binding in major depressive disorder after psychotherapy, but not after SSRI pharmacotherapy, is related to improved social functioning capacity. *Psychother Psychosom*. 2013;82(4):260-1. PMID: 23736831. Exclusion Code: X6

180. Kasper S, Gastpar M, Moller HJ, et al. Better tolerability of St. John's wort extract WS 5570 compared to treatment with SSRIs: a reanalysis of data from controlled clinical trials in acute major depression. *Int Clin Psychopharmacol*. 2010 Jul;25(4):204-13. PMID: 20568656. Exclusion Code: X4
181. Katon W, Robinson P, Von Korff M, et al. A multifaceted intervention to improve treatment of depression in primary care. *Arch Gen Psychiatry*. 1996 Oct;53(10):924-32. PMID: 8857869. Exclusion Code: X3
182. Katona C. New antidepressants for older people: a critical review of the evidence base. *Encephale*. 2008 Apr;34 Suppl 2:S71-S6. PMID: 18675006. Exclusion Code: X1
183. Katz T, Fisher P, Katz A, et al. The feasibility of a randomised, placebo-controlled clinical trial of homeopathic treatment of depression in general practice. *Homeopathy*; 2005. p. 145-52. Exclusion Code: X3
184. Keller MB, McCullough JP, Klein DN, et al. A comparison of nefazodone, the cognitive behavioral-analysis system of psychotherapy, and their combination for the treatment of chronic depression. *N Engl J Med*. 2000 May 18;342(20):1462-70. PMID: 10816183. Exclusion Code: X2
185. Kennedy SH, Segal ZV, Cohen NL, et al. Lithium carbonate versus cognitive therapy as sequential combination treatment strategies in partial responders to antidepressant medication: an exploratory trial. *J Clin Psychiatry*. 2003 Apr;64(4):439-44. PMID: 12716247. Exclusion Code: X2
186. Kessler D, Lewis G, Kaur S, et al. Therapist-delivered internet psychotherapy for depression in primary care: A randomised controlled trial. *The Lancet*. 2009;374(9690):628-34. PMID: 2009-18405-013. First Author & Affiliation: Kessler, David. Exclusion Code: X5
187. Kingston T, Dooley B, Bates A, et al. Mindfulness-based cognitive therapy for residual depressive symptoms. *Psychol Psychother*. 2007 Jun;80(Pt 2):193-203. PMID: 17535594. Exclusion Code: X4
188. Klainin-Yobas P, Cho MAA, Creedy D. Efficacy of mindfulness-based interventions on depressive symptoms among people with mental disorders: A meta-analysis. *Int J Nurs Stud*. 2012;49(1):109-21. PMID: 2012-01598-015. PMID: 21963234. First Author & Affiliation: Klainin-Yobas, Piyanee. Exclusion Code: X2
189. Kloiber S. Acupuncture for the treatment of major depressive disorder: A randomized controlled trial. *Deutsche zeitschrift fur akupunktur*; 2013. p. 35-6. Exclusion Code: X1
190. Kocsis JH, Gelenberg AJ, Rothbaum BO, et al. Cognitive behavioral analysis system of psychotherapy and brief supportive psychotherapy for augmentation of antidepressant nonresponse in chronic depression: the REVAMP Trial. *Arch Gen Psychiatry*. 2009 Nov;66(11):1178-88. PMID: 19884606. Exclusion Code: X7
191. Kocsis JH, Leon AC, Markowitz JC, et al. Patient preference as a moderator of outcome for chronic forms of major depressive disorder treated with nefazodone, cognitive behavioral analysis system of psychotherapy, or their combination. *J Clin Psychiatry*. 2009 Mar;70(3):354-61. PMID: 19192474. Exclusion Code: X2
192. Kocsis JH, Rush AJ, Markowitz JC, et al. Continuation treatment of chronic depression: a comparison of nefazodone, cognitive behavioral analysis system of psychotherapy, and their combination. *Psychopharmacol Bull*. 2003;37(4):73-87. PMID: 15131518. Exclusion Code: X2
193. Komossa K, Depping AM, Gaudchau A, et al. Second-generation antipsychotics for major depressive disorder and dysthymia. *Cochrane Database Syst Rev*. 2010(12):CD008121. PMID: 21154393. Exclusion Code: X3
194. Kramer U, de Roten Y, Perry JC, et al. Change in defense mechanisms and coping patterns during the course of 2-year-long psychotherapy and psychoanalysis for recurrent depression: a pilot study of a randomized controlled trial. *J Nerv Ment Dis*. 2013 Jul;201(7):614-20. PMID: 23817160. Exclusion Code: X5

195. Krishna M, Jauhari A, Lepping P, et al. Is group psychotherapy effective in older adults with depression? A systematic review. *Int J Geriatr Psychiatry*. 2011;26(4):331-40. PMID: 2011-07526-001. PMID: 20973096. First Author & Affiliation: Krishna, Murali. Exclusion Code: X5
196. Kriston L, von Wolff A, Westphal A, et al. EFFICACY AND ACCEPTABILITY OF ACUTE TREATMENTS FOR PERSISTENT DEPRESSIVE DISORDER: A NETWORK META-ANALYSIS. *Depress Anxiety*. 2014 Jan 21PMID: 24448972. Exclusion Code: X2
197. Krogh J, Nordentoft M, Sterne JAC, et al. The effect of exercise in clinically depressed adults: Systematic review and meta-analysis of randomized controlled trials. *J Clin Psychiatry*. 2011;72(4):529-38. PMID: 21034688. Exclusion Code: X3
198. Kronstrom K, Salminen JK, Hietala J, et al. Personality traits and recovery from major depressive disorder. *Nord J Psychiatry*. 2011 Feb;65(1):52-7. PMID: 20500120. Exclusion Code: X7
199. Kruisdijk FR, Hendriksen IJ, Tak EC, et al. Effect of running therapy on depression (EFFORT-D). Design of a randomised controlled trial in adult patients [ISRCTN 1894]. *BMC Public Health*. 2012;12:50. PMID: 22260713. Exclusion Code: X1
200. Lafferman J, Solomon K, Ruskin P. Lithium augmentation for treatment-resistant depression in the elderly. *J Geriatr Psychiatry Neurol*. 1988 Jan;1(1):49-52. PMID: 3150926. Exclusion Code: X4
201. Laidlaw K, Davidson K, Toner H, et al. A randomised controlled trial of cognitive behaviour therapy vs treatment as usual in the treatment of mild to moderate late life depression. *Int J Geriatr Psychiatry*. 2008 Aug;23(8):843-50. PMID: 18311844. Exclusion Code: X5
202. Lam RW, Kennedy SH. Evidence-based strategies for achieving and sustaining full remission in depression: focus on metaanalyses. *Can J Psychiatry*. 2004 Mar;49(3 Suppl 1):17S-26S. PMID: 15147033. Exclusion Code: X4
203. Lampe L, Coulston CM, Berk L. Psychological management of unipolar depression. *Acta Psychiatr Scand Suppl*. 2013(443):24-37. PMID: 23586874. Exclusion Code: X1
204. Landén M, Björling G, Agren H, et al. A randomized, double-blind, placebo-controlled trial of buspirone in combination with an SSRI in patients with treatment-refractory depression. *J Clin Psychiatry*; 1998. p. 664-8. Exclusion Code: X5
205. Lavretsky H. Combination of methylpehndate with citalopram is superior to either drug alone in improving clinical and cognitive outcomes in geriatric depression. *Neuropsychopharmacology*. 2013;38:S90. Exclusion Code: X9
206. Lavretsky H, Alstein LL, Olmstead RE, et al. Complementary use of tai chi chih augments escitalopram treatment of geriatric depression: a randomized controlled trial. *Am J Geriatr Psychiatry*. 2011 Oct;19(10):839-50. PMID: 21358389. Exclusion Code: X5
207. Lawlor DA, Hopker SW. The effectiveness of exercise as an intervention in the management of depression: Systematic review and meta-regression analysis of randomised controlled trials. *BMJ: British Medical Journal*. 2001;322(7289):763-6. PMID: 2001-17117-001. First Author & Affiliation: Lawlor, Debbie A. Exclusion Code: X5
208. Leo RJ, Ligot JSA, Jr. A systematic review of randomized controlled trials of acupuncture in the treatment of depression. *J Affect Disord*. 2007;97(1-3):13-22. PMID: 2006-23532-003. PMID: 16899301. First Author & Affiliation: Leo, Raphael J. Exclusion Code: X5
209. Lesperance F, Frasura-Smith N, St-Andre E, et al. The efficacy of omega-3 supplementation for major depression: a randomized controlled trial. *J Clin Psychiatry*. 2011 Aug;72(8):1054-62. PMID: 20584525. Exclusion Code: X2
210. Lester H, Gilbody S. Choosing a second generation antidepressant for treatment of major depressive disorder. *BMJ*. 2012;344:e1014. PMID: 22334542. Exclusion Code: X1

211. Lewis AJ, Dennerstein M, Gibbs PM. Short-term psychodynamic psychotherapy: Review of recent process and outcome studies. *Aust N Z J Psychiatry*. 2008;42(6):445-55. PMID: 2008-12696-002. PMID: 18465371. First Author & Affiliation: Lewis, Andrew J. Exclusion Code: X2
212. Lin C-H, Lin S-H, Jang F-L. Adjunctive low-dose aripiprazole with standard-dose sertraline in treating fresh major depressive disorder. *J Clin Psychopharmacol*. 2011;31(5):563-8. PMID: 2011-19925-003. PMID: 21869699. First Author & Affiliation: Lin, Chien-Ho. Exclusion Code: X5
213. Ludman EJ, Simon GE, Tutty S, et al. A randomized trial of telephone psychotherapy and pharmacotherapy for depression: continuation and durability of effects. *J Consult Clin Psychol*. 2007 Apr;75(2):257-66. PMID: 17469883. Exclusion Code: X5
214. Lynch TR, Morse JQ, Mendelson T, et al. Dialectical behavior therapy for depressed older adults: A randomized pilot study. *The American Journal of Geriatric Psychiatry*. 2003;11(1):33-45. PMID: 2003-01261-008. PMID: 12527538. First Author & Affiliation: Lynch, Thomas R. Exclusion Code: X5
215. Lyons Z, van dWG, Shen Z, et al. Acupuncture and Chinese herbs as treatments for depression: An Australian pilot study. *Complement Ther Clin Pract*. 2012 Nov;18(4):216-20. PMID: 0165371. Exclusion Code: X3
216. Ma SH, Teasdale JD. Mindfulness-based cognitive therapy for depression: replication and exploration of differential relapse prevention effects. *J Consult Clin Psychol*. 2004 Feb;72(1):31-40. PMID: 14756612. Exclusion Code: X2
217. Malik ML, Beutler LE, Alimohamed S, et al. Are all cognitive therapies alike? A comparison of cognitive and noncognitive therapy process and implications for the application of empirically supported treatments. *J Consult Clin Psychol*. 2003 Feb;71(1):150-8. PMID: 12602435. Exclusion Code: X4
218. Manber R, Arnow B, Blasey C, et al. Patient's therapeutic skill acquisition and response to psychotherapy, alone or in combination with medication. *Psychol Med*. 2003 May;33(4):693-702. PMID: 12785471. Exclusion Code: X6
219. Manber R, Edinger JD, Gress JL, et al. Cognitive behavioral therapy for insomnia enhances depression outcome in patients with comorbid major depressive disorder and insomnia. *Sleep*. 2008 Apr;31(4):489-95. PMID: 18457236. Exclusion Code: X2
220. Manber R, Rush AJ, Thase ME, et al. The effects of psychotherapy, nefazodone, and their combination on subjective assessment of disturbed sleep in chronic depression. *Sleep*. 2003 Mar 15;26(2):130-6. PMID: 12683470. Exclusion Code: X2
221. Maron E, Eller T, Vasar V, et al. Effects of bupropion augmentation in escitalopram-resistant patients with major depressive disorder: An open-label, naturalistic study. *J Clin Psychiatry*. 2009;70(7):1054-6. Exclusion Code: X5
222. Martin S. A randomised controlled trial of cognitive therapy versus anti-depressants for major depression with sequential SPECT scanning to measure changes in cerebral blood flow with treatment. *National Research Register*; 1999. Exclusion Code: X9
223. Martin SD, Martin E, Rai SS, et al. Brain blood flow changes in depressed patients treated with interpersonal psychotherapy or venlafaxine hydrochloride: preliminary findings. *Arch Gen Psychiatry*. 2001 Jul;58(7):641-8. PMID: 11448369. Exclusion Code: X4
224. Martiny K, Lunde M, Bech P, et al. A short-term double-blind randomized controlled pilot trial with active or placebo pindolol in patients treated with venlafaxine for major depression. *Nord J Psychiatry*. 2012 Jun;66(3):147-54. PMID: 22458638. Exclusion Code: X2

225. Mather AS, Rodriguez C, Guthrie MF, et al. Effects of exercise on depressive symptoms in older adults with poorly responsive depressive disorder: Randomised controlled trial. *Br J Psychiatry*. 2002;180(5):411-5. PMID: 2002-13169-003. First Author & Affiliation: Mather, Anne S. Exclusion Code: X5
226. Matreja PS, Badyal DK, Deswal RS, et al. Efficacy and safety of add on low-dose mirtazapine in depression. *Indian J Pharmacol*. 2012;44(2):173-7. Exclusion Code: X2
227. Mazzucchelli T, Kane R, Rees C. Behavioral activation treatments for depression in adults: A meta-analysis and review. *Clinical Psychology: Science and Practice*. 2009;16(4):383-411. PMID: 2009-20281-001. First Author & Affiliation: Mazzucchelli, Trevor. Exclusion Code: X8
228. McBride C, Segal Z, Kennedy S, et al. Changes in autobiographical memory specificity following cognitive behavior therapy and pharmacotherapy for major depression. *Psychopathology*. 2007;40(3):147-52. PMID: 17318006. Exclusion Code: X5
229. McIntyre A, Gendron A, McIntyre A. Quetiapine adjunct to selective serotonin reuptake inhibitors or venlafaxine in patients with major depression, comorbid anxiety, and residual depressive symptoms: a randomized, placebo-controlled pilot study. *Depress Anxiety*. 2007;24(7):487-94. PMID: 17177199. Exclusion Code: X5
230. McPherson S, Cairns P, Carlyle J, et al. The effectiveness of psychological treatments for treatment-resistant depression: a systematic review. *Acta Psychiatr Scand*. 2005 May;111(5):331-40. PMID: 15819726. Exclusion Code: X4
231. Mehta P, Sharma M. Yoga as a complementary therapy for clinical depression. *Complementary Health Practice Review*. 2010;15(3):156-70. PMID: 2010-25846-004. First Author & Affiliation: Mehta, Purvi. Exclusion Code: X1
232. Mergl R, Henkel V, Allgaier A-K, et al. Are treatment preferences relevant in response to serotonergic antidepressants and cognitive-behavioral therapy in depressed primary care patients? Results from a randomized controlled trial including a patients' choice arm. *Psychother Psychosom*. 2010;80(1):39-47. PMID: 2010-26489-005. PMID: 20975325. First Author & Affiliation: Mergl, Roland. Exclusion Code: X2
233. Minelli A, Zambello F, Vaona A. Efficacia della psicoterapia cognitivo-comportamentale associata a terapia psicofarmacologica nella depressione. Revisione della letteratura metanalitica. *Rivista di Psichiatria*. 2011;46(1):18-23. PMID: 2011-05487-002. PMID: 21446108. First Author & Affiliation: Minelli, Alessandra. Exclusion Code: X1
234. Miniati M, Rucci P, Frank E, et al. Sensitivity to change and predictive validity of the MOODS-SR questionnaire, last-month version. *Psychother Psychosom*. 2009;78(2):116-24. PMID: 19218830. Exclusion Code: X5
235. Miranda J, Green BL, Krupnick JL, et al. One-year outcomes of a randomized clinical trial treating depression in low-income minority women. *J Consult Clin Psychol*. 2006;74(1):99-111. PMID: 2006-03253-010. PMID: 16551147. First Author & Affiliation: Miranda, Jeanne. Exclusion Code: X6
236. Miser WF. Exercise as an effective treatment option for major depression in older adults. *J Fam Pract*. 2000 Feb;49(2):109-10. PMID: 10718684. Exclusion Code: X1
237. Mohr DC, Carmody T, Erickson L, et al. Telephone-administered cognitive behavioral therapy for veterans served by community-based outpatient clinics. *J Consult Clin Psychol*. 2011 Apr;79(2):261-5. PMID: 21299274. Exclusion Code: X5
238. Molenaar PJ, Dekker J, Van R, et al. Does adding psychotherapy to pharmacotherapy improve social functioning in the treatment of outpatient depression? *Depress Anxiety*. 2007;24(8):553-62. PMID: 2007-19727-004. PMID: 17131302. First Author & Affiliation: Molenaar, Pieter J. Exclusion Code: X5

239. Moller H. Johanniskraut in der primär-ärztlichen Therapie: Hypericum als fester Bestandteil in der Therapie von Depressionen. *Schweiz Z Ganzheits Medizin*. 2004 Nov;16(7-8):391-2. PMID: 0082428. Exclusion Code: X1
240. Moore RG, Blackburn IM. Cognitive therapy in the treatment of non-responders to antidepressant medication: A controlled pilot study. *Behav Cogn Psychother*; 1997. p. 251-9. Exclusion Code: X5
241. Moradveisi L, Huibers M, Renner F, et al. The influence of patients' preference/attitude towards psychotherapy and antidepressant medication on the treatment of major depressive disorder. *J Behav Ther Exp Psychiatry*. 2014;45(1):170-7. Exclusion Code: X4
242. Mota-Pereira J, Silverio J, Carvalho S, et al. Moderate exercise improves depression parameters in treatment-resistant patients with major depressive disorder. *J Psychiatr Res*. 2011 Aug;45(8):1005-11. PMID: 21377690. Exclusion Code: X2
243. Mozaffari-Khosravi H, Yassini-Ardakani M, Karamati M, et al. Eicosapentaenoic acid versus docosahexaenoic acid in mild-to-moderate depression: A randomized, double-blind, placebo-controlled trial. *Eur Neuropsychopharmacol*. 2013;23(7):636-44. PMID: 2012-22973-001. PMID: 22910528. First Author & Affiliation: Mozaffari-Khosravi, Hassan. Exclusion Code: X2
244. Mukaino Y, Park J, White A, et al. The effectiveness of acupuncture for depression - a systematic review of randomised controlled trials. *Acupunct Med*. 2005 Jun;23(2):70-6. PMID: 0075492. Exclusion Code: X5
245. Mulrow CD, Williams JW, Jr., Trivedi M, et al. Treatment of depression--newer pharmacotherapies. *Psychopharmacol Bull*. 1998;34(4):409-795. PMID: 10513454. Exclusion Code: X1
246. Mura G, Moro MF, Patten SB, et al. Exercise as an add-on strategy for the treatment of major depressive disorder: a systematic review. *CNS Spectr*. 2014 Mar 3;1-13. PMID: 24589012. Exclusion Code: X3
247. Mynors-Wallis L. Problem-solving treatment: evidence for effectiveness and feasibility in primary care. *Int J Psychiatry Med*. 1996;26(3):249-62. PMID: 8976466. Exclusion Code: X4
248. Nelson JC. S-adenosyl methionine (SAmE) augmentation in major depressive disorder. *Am J Psychiatry*. 2010 Aug;167(8):889-91. PMID: 20693465. Exclusion Code: X1
249. Nelson MK. Meta-analysis: Hypnotherapy/cognitive-behavioral therapy and its efficacy on depression compared to pharmacotherapy. US: ProQuest Information & Learning; 2002. Exclusion Code: X1
250. Nemeroff CB, Amchin J. Placebo-controlled trial of the efficacy and tolerability of venlafaxine and fluoxetine in outpatients with major depression. 11th European College of Neuropsychopharmacology Congress. Paris, France. 31st October 4th November 1998.; 1998. Exclusion Code: X9
251. Nemets B, Stahl Z, Belmaker RH. Addition of omega-3 fatty acid to maintenance medication treatment for recurrent unipolar depressive disorder. *Am J Psychiatry*. 2002 Mar;159(3):477-9. PMID: 11870016. Exclusion Code: X2
252. Newhouse PA. Use of serotonin selective reuptake inhibitors in geriatric depression. *J Clin Psychiatry*. 1996;57 Suppl 5:12-22. PMID: 8647788. Exclusion Code: X1
253. Nieuwsma JA, Trivedi RB, McDuffie J, et al. Brief psychotherapy for depression: A systematic review and meta-analysis. *Int J Psychiatry Med*. 2012;43(2):129-51. PMID: 2012-17351-003. PMID: 22849036. First Author & Affiliation: Nieuwsma, Jason A. Exclusion Code: X5
254. Ninan PT, Rush AJ, Crits-Christoph P, et al. Symptomatic and syndromal anxiety in chronic forms of major depression: effect of nefazodone, cognitive behavioral analysis system of psychotherapy, and their combination. *J Clin Psychiatry*. 2002 May;63(5):434-41. PMID: 12025827. Exclusion Code: X3

255. Oeland A-M, Laessoe U, Olesen AV, et al. Impact of exercise on patients with depression and anxiety. *Nordic Journal of Psychiatry*. 2010;64(3):210-7. PMID: 2010-09396-010. PMID: 20100135. First Author & Affiliation: Oeland, Anne-Marie. Exclusion Code: X2
256. Onder E, Tural U. Faster response in depressive patients treated with fluoxetine alone than in combination with buspirone. *J Affect Disord*. 2003 Sep;76(1-3):223-7. PMID: 12943952. Exclusion Code: X5
257. Otto M, Wisniewski S. CBT for treatment resistant depression. *Lancet*. 2013 Feb 2;381(9864):352-3. PMID: 0165503. Exclusion Code: X1
258. Pae CU, Patkar AA, Jun TY, et al. Aripiprazole augmentation for treatment of patients with inadequate antidepressants response. *Depress Anxiety*. 2007;24(7):522-6. Exclusion Code: X5
259. Pampallona S, Bollini P, Tibaldi G, et al. Combined pharmacotherapy and psychological treatment for depression: a systematic review. *Arch Gen Psychiatry*. 2004 Jul;61(7):714-9. PMID: 15237083. Exclusion Code: X5
260. Papakostas GI, Cassiello CF, Iovieno N. Folate and S-adenosylmethionine for major depressive disorder. *Can J Psychiatry*. 2012 Jul;57(7):406-13. PMID: 22762295. Exclusion Code: X1
261. Papakostas GI, Fava M, Thase ME. Treatment of SSRI-resistant depression: a meta-analysis comparing within- versus across-class switches. *Biol Psychiatry*. 2008 Apr 1;63(7):699-704. PMID: 17919460. Exclusion Code: X2
262. Papakostas GI, Mischoulon D, Shyu I, et al. S-adenosyl methionine (S-AdMet) augmentation of serotonin reuptake inhibitors for antidepressant nonresponders with major depressive disorder: a double-blind, randomized clinical trial. *Am J Psychiatry*. 2010 Aug;167(8):942-8. PMID: 20595412. Exclusion Code: X2
263. Papakostas GI, Petersen TJ, Kinrys G, et al. Aripiprazole augmentation of selective serotonin reuptake inhibitors for treatment-resistant major depressive disorder. *J Clin Psychiatry*. 2005;66(10):1326-30. Exclusion Code: X5
264. Parker G, Brotchie H, Parker K. Is combination olanzapine and antidepressant medication associated with a more rapid response trajectory than antidepressant alone? *Am J Psychiatry*. 2005;162(4):796-8. Exclusion Code: X5
265. Parker G, Paterson A, Blanch B. Suggested early onset of true action of antidepressant drugs may be artefactual: a heuristic study. *Int Clin Psychopharmacol*. 2013 Jan;28(1):29-32. PMID: 23232755. Exclusion Code: X2
266. Parker G, Roy K, Wilhelm K, et al. Assessing the comparative effectiveness of antidepressant therapies: a prospective clinical practice study. *J Clin Psychiatry*. 2001 Feb;62(2):117-25. PMID: 11247097. Exclusion Code: X2
267. Patkar AA, Masand PS, Pae CU, et al. A randomized, double-blind, placebo-controlled trial of augmentation with an extended release formulation of methylphenidate in outpatients with treatment-resistant depression. *J Clin Psychopharmacol*. 2006;26(6):653-6. Exclusion Code: X5
268. Paykel ES, Scott J, Cornwall PL, et al. Duration of relapse prevention after cognitive therapy in residual depression: follow-up of controlled trial. *Psychol Med*. 2005 Jan;35(1):59-68. PMID: 15842029. Exclusion Code: X1
269. Paykel ES, Scott J, Teasdale JD, et al. Prevention of relapse in residual depression by cognitive therapy: a controlled trial. *Arch Gen Psychiatry*. 1999 Sep;56(9):829-35. PMID: 12884889. Exclusion Code: X3
270. Peet M, Horrobin DF. A dose-ranging study of the effects of ethyl-eicosapentaenoate in patients with ongoing depression despite apparently adequate treatment with standard drugs. *Arch Gen Psychiatry*. 2002 Oct;59(10):913-9. PMID: 12365878. Exclusion Code: X2
271. Pereira D, de QB, Miranda A, et al. Effects of physical exercise on plasma levels of brain-derived neurotrophic factor and depressive symptoms in elderly women-a randomized clinical trial. *Arch Phys Med Rehabil*. 2013 Aug;94(8):1443-50. PMID: 0170573. Exclusion Code: X5

272. Perez V, Puigdemont D, Gilaberte I, et al. Augmentation of fluoxetine's antidepressant action by pindolol: analysis of clinical, pharmacokinetic, and methodologic factors. *J Clin Psychopharmacol*. 2001 Feb;21(1):36-45. PMID: 11199945. Exclusion Code: X5
273. Perez V, Soler J, Puigdemont D, et al. A double-blind, randomized, placebo-controlled trial of pindolol augmentation in depressive patients resistant to serotonin reuptake inhibitors. *Grup de Recerca en Trastorns Afectius. Arch Gen Psychiatry*. 1999 Apr;56(4):375-9. PMID: 10197835. Exclusion Code: X2
274. Perlis RH, Alpert J, Nierenberg AA, et al. Clinical and sociodemographic predictors of response to augmentation, or dose increase among depressed outpatients resistant to fluoxetine 20 mg/day. *Acta Psychiatr Scand*. 2003 Dec;108(6):432-8. PMID: 14616224. Exclusion Code: X5
275. Perlis RH, Nierenberg AA, Alpert JE, et al. Effects of adding cognitive therapy to fluoxetine dose increase on risk of relapse and residual depressive symptoms in continuation treatment of major depressive disorder. *J Clin Psychopharmacol*. 2002 Oct;22(5):474-80. PMID: 12352270. Exclusion Code: X2
276. Perry EB, Berman RM, Sanacora G, et al. Pindolol augmentation in depressed patients resistant to selective serotonin reuptake inhibitors: a double-blind, randomized, controlled trial. *J Clin Psychiatry*. 2004 Feb;65(2):238-43. PMID: 15003079. Exclusion Code: X5
277. Petersen T, Harley R, Papakostas GI, et al. Continuation cognitive-behavioural therapy maintains attributional style improvement in depressed patients responding acutely to fluoxetine. *Psychol Med*. 2004 Apr;34(3):555-61. PMID: 15259840. Exclusion Code: X2
278. Peterson TJ, Feldman G, Harley R, et al. Extreme response style in recurrent and chronically depressed patients: change with antidepressant administration and stability during continuation treatment. *J Consult Clin Psychol*. 2007 Feb;75(1):145-53. PMID: 17295573. Exclusion Code: X2
279. Pilkington K, Kirkwood G, Rampes H, et al. Yoga for depression: The research evidence. *J Affect Disord*. 2005;89(1-3):13-24. PMID: 2006-00510-002. PMID: 16185770. First Author & Affiliation: Pilkington, Karen. Exclusion Code: X5
280. Pinquart M, Duberstein PR, Lyness JM. Treatments for later-life depressive conditions: a meta-analytic comparison of pharmacotherapy and psychotherapy. *Am J Psychiatry*. 2006 Sep;163(9):1493-501. PMID: 16946172. Exclusion Code: X2
281. Pinquart M, Duberstein PR, Lyness JM. Effects of pharmacotherapy and psychotherapy in late-life depression. *Brown University Psychopharmacology Update*. 2006;17(12):1. PMID: 2009361051. Language: English. Entry Date: 20080125. Publication Type: journal article. Journal Subset: Biomedical. Exclusion Code: X4
282. Posternak M, Novak S, Stern R, et al. A pilot effectiveness study: placebo-controlled trial of adjunctive L-triiodothyronine (T3) used to accelerate and potentiate the antidepressant response. *Int J Neuropsychopharmacol*. 2008 Feb;11(1):15-25. PMID: 17352847. Exclusion Code: X2
283. Potter C. Depression - psychosoziale Selbsttötung: Selbstbehauptungsstörungen als Ausdruck einer tiefen Identitätskrise durch Adaptation falscher Verhaltensmuster. *Naturheilkunde*. 2006;2006(5):22-4. PMID: 0093045. Exclusion Code: X1
284. Proudfoot J, Goldberg D, Mann A, et al. Computerized, interactive, multimedia cognitive-behavioural program for anxiety and depression in general practice. *Psychol Med*. 2003 Feb;33(2):217-27. PMID: 12622301. Exclusion Code: X2
285. Proudfoot J, Ryden C, Everitt B, et al. Clinical efficacy of computerised cognitive-behavioural therapy for anxiety and depression in primary care: Randomised controlled trial. *Br J Psychiatry*. 2004;185(1):46-54. PMID: 2004-16407-010. First Author & Affiliation: Proudfoot, Judith. Exclusion Code: X5

286. Quilty LC, De Fruyt F, Rolland JP, et al. Dimensional personality traits and treatment outcome in patients with major depressive disorder. *J Affect Disord.* 2008 Jun;108(3):241-50. PMID: 18067975. Exclusion Code: X4
287. Rafanelli C, Ruini C, Belaise C, et al. Approccio cognitive comportamentale alla perdita di efficacia durante la terapia di mantenimento con antidepressivi. *Rivista di Psichiatria.* 2003;38(5):259-62. PMID: 2003-10067-004. First Author & Affiliation: Rafanelli, Chiara. Exclusion Code: X5
288. Rahimi R, Nikfar S, Abdollahi M. Efficacy and tolerability of *Hypericum perforatum* in major depressive disorder in comparison with selective serotonin reuptake inhibitors: a meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry.* 2009 Feb 1;33(1):118-27. PMID: 19028540. Exclusion Code: X8
289. Rapaport MH, Gharabawi GM, Canuso CM, et al. Corrigendum: Effects of Risperidone Augmentation in Patients with Treatment-Resistant Depression: Results of Open-Label Treatment Followed by Double-Blind Continuation. *Neuropsychopharmacology.* 2006;31(11) PMID: 2006-20317-023. First Author & Affiliation: Rapaport, Mark Hyman. Exclusion Code: X1
290. Ravindran AV, Kennedy SH, O'Donovan MC, et al. Osmotic-release oral system methylphenidate augmentation of antidepressant monotherapy in major depressive disorder: Results of a double-blind, randomized, placebo-controlled trial. *J Clin Psychiatry.* 2008;69(1):87-94. Exclusion Code: X5
291. Reeves H, Batra S, May RS, et al. Efficacy of risperidone augmentation to antidepressants in the management of suicidality in major depressive disorder: A randomized, double-blind, placebo-controlled pilot study. *J Clin Psychiatry.* 2008;69(8):1228-36. Exclusion Code: X5
292. Rethorst CD, Wipfli BM, Landers DM. The antidepressive effects of exercise: A meta-analysis of randomized trials. *Sports Med.* 2009;39(6):491-511. PMID: 2009-08145-001. PMID: 19453207. First Author & Affiliation: Rethorst, Chad D. Exclusion Code: X5
293. Reynolds CF, III, Dew MA, Martire LM, et al. Treating depression to remission in older adults: A controlled evaluation of combined escitalopram with interpersonal psychotherapy versus escitalopram with depression care management. *Int J Geriatr Psychiatry.* 2010;25(11):1134-41. PMID: 2010-23017-006. PMID: 20957693. First Author & Affiliation: Reynolds, Charles F. III. Exclusion Code: X5
294. Robertson R, Robertson A, Jepson R, et al. Walking for depression or depressive symptoms: A systematic review and meta-analysis. *Mental Health and Physical Activity.* 2012;5(1):66-75. Exclusion Code: X2
295. Röder C, Schaefer M, Leucht S. Meta-Analyse zu Wirksamkeit und Verträglichkeit der Behandlung der leichten und mittelschweren Depression mit Johanniskraut. *Fortschritte der Neurologie, Psychiatrie.* 2004;72(6):330-43. PMID: 2004-15951-002. First Author & Affiliation: Röder, Claudia. Exclusion Code: X4
296. Roshanaei-Moghaddam B, Pauly MC, Atkins DC, et al. Relative effects of CBT and pharmacotherapy in depression versus anxiety: is medication somewhat better for depression, and CBT somewhat better for anxiety? *Depress Anxiety.* 2011 Jul;28(7):560-7. PMID: 21608087. Exclusion Code: X2
297. Ruhe HG, Huyser J, Swinkels JA, et al. Switching antidepressants after a first selective serotonin reuptake inhibitor in major depressive disorder: a systematic review. *J Clin Psychiatry.* 2006 Dec;67(12):1836-55. PMID: 17194261. Exclusion Code: X2
298. Rush AJ. Combining antidepressant medications: a good idea? *Am J Psychiatry.* 2010 Mar;167(3):241-3. PMID: 20194484. Exclusion Code: X1
299. Rush AJ, Fava M, Wisniewski SR, et al. Sequenced treatment alternatives to relieve depression (STAR*D): rationale and design. *Control Clin Trials.* 2004 Feb;25(1):119-42. PMID: 15061154. Exclusion Code: X4

300. Rush AJ, Trivedi MH, Carmody TJ, et al. Self-reported depressive symptom measures: Sensitivity to detecting change in a randomized, controlled trial of chronically depressed, nonpsychotic outpatients. *Neuropsychopharmacology*. 2005;30(2):405-16. Exclusion Code: X4
301. Rush AJ, Wisniewski SR, Warden D, et al. Selecting among second-step antidepressant medication monotherapies: predictive value of clinical, demographic, or first-step treatment features. *Arch Gen Psychiatry*. 2008 Aug;65(8):870-80. PMID: 18678792. Exclusion Code: X4
302. Rutherford B, Sneed J, Miyazaki M, et al. An open trial of aripiprazole augmentation for SSRI non-remitters with late-life depression. *Int J Geriatr Psychiatry*. 2007;22(10):986-91. Exclusion Code: X5
303. Saiz-Ruiz J, Ibaez A, Diaz-Marsa M, et al. Efficacy of venlafaxine in major depression resistant to selective serotonin reuptake inhibitors. *Prog Neuropsychopharmacol Biol Psychiatry*. 2002;26(6):1129-34. Exclusion Code: X5
304. Samad Z, Brealey S, Gilbody S. The effectiveness of behavioural therapy for the treatment of depression in older adults: A meta-analysis. *Int J Geriatr Psychiatry*. 2011;26(12):1211-20. PMID: 2012-11062-001. PMID: 21308789. First Author & Affiliation: Samad, Zara. Exclusion Code: X5
305. Sarris J, Fava M, Schweitzer I, et al. St John's wort (*Hypericum perforatum*) versus sertraline and placebo in major depressive disorder: continuation data from a 26-week RCT. *Pharmacopsychiatry*. 2012 Nov;45(7):275-8. PMID: 22592504. Exclusion Code: X4
306. Sarris J, Schoendorfer N, Kavanagh DJ. Major depressive disorder and nutritional medicine: a review of monotherapies and adjuvant treatments. *Nutr Rev*. 2009 Mar;67(3):125-31. PMID: 19239627. Exclusion Code: X8
307. Schaefer KL. Residual symptoms after treatment of chronic depression: A comparison across treatment modalities. US: ProQuest Information & Learning; 2008. Exclusion Code: X2
308. Schatzberg AF, Rush AJ, Arnow BA, et al. Chronic depression: medication (nefazodone) or psychotherapy (CBASP) is effective when the other is not. *Arch Gen Psychiatry*. 2005 May;62(5):513-20. PMID: 15867104. Exclusion Code: X4
309. Schuch FB, Vasconcelos-Moreno MP, Fleck MP. The impact of exercise on quality of life within exercise and depression trials: A systematic review. *Mental Health and Physical Activity*. 2011;4(2):43-8. Exclusion Code: X2
310. Schwartz TL, Nasra GS, Chilton M, et al. Aripiprazole augmentation of selective serotonin or serotonin norepinephrine reuptake inhibitors in the treatment of major depressive disorder. *Primary Psychiatry*. 2007;14(1):67-9. Exclusion Code: X5
311. Scott C, Tacchi MJ, Jones R, et al. Acute and one-year outcome of a randomised controlled trial of brief cognitive therapy for major depressive disorder in primary care. *Br J Psychiatry*. 1997 Aug;171:131-4. PMID: 9337947. Exclusion Code: X5
312. Scott J, Palmer S, Paykel E, et al. Use of cognitive therapy for relapse prevention in chronic depression. Cost-effectiveness study. *Br J Psychiatry*. 2003 Mar;182:221-7. PMID: 12611785. Exclusion Code: X2
313. Scott J, Teasdale JD, Paykel ES, et al. Effects of cognitive therapy on psychological symptoms and social functioning in residual depression. *Br J Psychiatry*. 2000 Nov;177:440-6. PMID: 11059998. Exclusion Code: X3
314. Segal Z, Vincent P, Levitt A. Efficacy of combined, sequential and crossover psychotherapy and pharmacotherapy in improving outcomes in depression. *J Psychiatry Neurosci*. 2002 Jul;27(4):281-90. PMID: 12174737. Exclusion Code: X1
315. Seo HJ, Jung YE, Woo YS, et al. Effect of augmented atypical antipsychotics on weight change in patients with major depressive disorder in a naturalistic setting. *Hum Psychopharmacol*. 2009 Mar;24(2):135-43. PMID: 19156709. Exclusion Code: X2

316. Serfaty MA, Haworth D, Blanchard M, et al. Clinical effectiveness of individual cognitive behavioral therapy for depressed older people in primary care: a randomized controlled trial. *Arch Gen Psychiatry*. 2009 Dec;66(12):1332-40. PMID: 19996038. Exclusion Code: X3
317. Sharma VK, Das S, Mondal S, et al. Effect of Sahaj Yoga on neuro-cognitive functions in patients suffering from major depression. *Indian J Physiol Pharmacol*. 2006 Oct-Dec;50(4):375-83. PMID: 17402267. Exclusion Code: X5
318. Sharma VK, Das S, Mondal S, et al. Effect of Sahaj Yoga on depressive disorders. *Indian J Physiol Pharmacol*. 2005 Oct-Dec;49(4):462-8. PMID: 16579401. Exclusion Code: X3
319. Shelton RC, Papakostas GI. Augmentation of antidepressants with atypical antipsychotics for treatment-resistant major depressive disorder. *Acta Psychiatr Scand*. 2008 Apr;117(4):253-9. PMID: 18190674. Exclusion Code: X4
320. Shelton RC, Tollefson GD, Tohen M, et al. A novel augmentation strategy for treating resistant major depression. *Am J Psychiatry*. 2001 Jan;158(1):131-4. PMID: 11136647. Exclusion Code: X2
321. Shimodera S, Watanabe N, Furukawa TA, et al. Change in quality of life after brief behavioral therapy for insomnia in concurrent depression: analysis of the effects of a randomized controlled trial. *J Clin Sleep Med*. 2014;10(4):433-9. PMID: 24733990. Exclusion Code: X3
322. Shirazi MM. St. John's wort doesn't help with major depression. *Tufts University Health & Nutrition Letter*. 2001;19(4):2-. Exclusion Code: X1
323. Siddique J, Chung JY, Brown CH, et al. Comparative effectiveness of medication versus cognitive-behavioral therapy in a randomized controlled trial of low-income young minority women with depression. *J Consult Clin Psychol*. 2012 Dec;80(6):995-1006. PMID: 23088620. Exclusion Code: X4
324. Sikorski C, Lupp M, Kersting A, et al. Effektivität computer- und internetgestützter kognitiver Verhaltenstherapie bei Depression: Ein systematischer Literaturüberblick. *Psychiatr Prax*. 2011;38(2):61-8. PMID: 2011-08849-003. PMID: 20972949. First Author & Affiliation: Sikorski, Claudia. Exclusion Code: X5
325. Simon JS, Nemeroff CB. Aripiprazole augmentation of antidepressants for the treatment of partially responding and nonresponding patients with major depressive disorder. *J Clin Psychiatry*. 2005 Oct;66(10):1216-20. PMID: 16259533. Exclusion Code: X5
326. Singh NA, Stavrinou TM, Scarbek Y, et al. A randomized controlled trial of high versus low intensity weight training versus general practitioner care for clinical depression in older adults. *J Gerontol A Biol Sci Med Sci*. 2005 Jun;60(6):768-76. PMID: 15983181. Exclusion Code: X2
327. Sjosten N, Kivela SL. The effects of physical exercise on depressive symptoms among the aged: a systematic review. *Int J Geriatr Psychiatry*. 2006 May;21(5):410-8. PMID: 16676285. Exclusion Code: X5
328. Smith A, Jain AK, Bennet C. Sertraline for recurrent major depression. *Am J Psychiatry*. 2005 May;162(5):1025-6. PMID: 15863821. Exclusion Code: X1
329. Sokolski KN. Adjunctive aripiprazole for bupropion-resistant major depression. *Ann Pharmacother*. 2008;42(7-8):1124-9. PMID: 2010001602. Language: English. Entry Date: 20080926. Publication Type: journal article. Exclusion Code: X1
330. Sokolski KN, Conney JC, Brown BJ, et al. Once-daily high-dose pindolol for SSRI-refractory depression. *Psychiatry Res*. 2004 Feb 15;125(2):81-6. PMID: 15006431. Exclusion Code: X2
331. Spek V, Cuijpers P, Nyklíček I, et al. Internet-based cognitive behaviour therapy for symptoms of depression and anxiety: A meta-analysis. *Psychological Medicine: A Journal of Research in Psychiatry and the Allied Sciences*. 2007;37(3):319-28. PMID: 2007-03729-002. First Author & Affiliation: Spek, Viola. Exclusion Code: X5

332. Spielmans GI, Berman MI, Linardatos E, et al. Adjunctive atypical antipsychotic treatment for major depressive disorder: a meta-analysis of depression, quality of life, and safety outcomes. *PLoS Med*. 2013;10(3):e1001403. PMID: 23554581. Exclusion Code: X5
333. Spielmans GI, Berman MI, Usitalo AN. Psychotherapy versus second-generation antidepressants in the treatment of depression: A meta-analysis. *J Nerv Ment Dis*. 2011;199(3):142-9. PMID: 21346483. Exclusion Code: X3
334. Spijker J, van Straten A, Bockting CL, et al. Psychotherapy, antidepressants, and their combination for chronic major depressive disorder: a systematic review. *Can J Psychiatry*. 2013 Jul;58(7):386-92. PMID: 23870720. Exclusion Code: X8
335. Steffens DC, Doraiswamy PM, McQuoid DR. Bupropion SR in the naturalistic treatment of elderly patients with major depression. *Int J Geriatr Psychiatry*. 2001;16(9):862-5. Exclusion Code: X7
336. Steidtmann D, Manber R, Arnow BA, et al. Patient treatment preference as a predictor of response and attrition in treatment for chronic depression. *Depress Anxiety*. 2012;29(10):896-905. Exclusion Code: X3
337. Stevinson C. Negative result for St John's wort in major depression. *Focus on Alternative and Complementary Therapies*. 2001 Sep;6(3):196-7. PMID: 0029317. Exclusion Code: X5
338. Stich FA. A meta-analysis of physical exercise as a treatment for symptoms of anxiety and depression. *US: ProQuest Information & Learning*; 1999. Exclusion Code: X5
339. Sturmey P. Behavioral activation is an evidence-based treatment for depression. *Behav Modif*. 2009;33(6):818-29. PMID: 2009-22984-006. PMID: 19933444. First Author & Affiliation: Sturmey, Peter. Exclusion Code: X5
340. Tadic A, Gorbulev S, Dahmen N, et al. Rationale and design of the randomised clinical trial comparing early medication change (EMC) strategy with treatment as usual (TAU) in patients with Major Depressive Disorder - the EMC trial. *Trials*. 2010;11. Exclusion Code: X3
341. Taneja C, Papakostas GI, Jing Y, et al. Cost-effectiveness of adjunctive therapy with atypical antipsychotics for acute treatment of major depressive disorder. *Ann Pharmacother*. 2012;46(5):642-9. PMID: 2011549176. Language: English. Entry Date: 20130614. Revision Date: 20130614. Publication Type: journal article. Exclusion Code: X1
342. Teasdale JD, Moore RG, Hayhurst H, et al. Metacognitive awareness and prevention of relapse in depression: empirical evidence. *J Consult Clin Psychol*. 2002 Apr;70(2):275-87. PMID: 11952186. Exclusion Code: X2
343. Thase ME, Friedman ES, Fasiczka AL, et al. Treatment of men with major depression: a comparison of sequential cohorts treated with either cognitive-behavioral therapy or newer generation antidepressants. *J Clin Psychiatry*. 2000 Jul;61(7):466-72. PMID: 10937603. Exclusion Code: X2
344. Thase ME, Greenhouse JB, Frank E, et al. Treatment of major depression with psychotherapy or psychotherapy-pharmacotherapy combinations. *Arch Gen Psychiatry*. 1997 Nov;54(11):1009-15. PMID: 9366657. Exclusion Code: X3
345. Thase ME, Rush AJ, Manber R, et al. Differential effects of nefazodone and cognitive behavioral analysis system of psychotherapy on insomnia associated with chronic forms of major depression. *J Clin Psychiatry*. 2002 Jun;63(6):493-500. PMID: 12088160. Exclusion Code: X4
346. Thomas P, Bordet R, Alexandre JY, et al. Pindolol addition shorten delay of action of paroxetine in major depression: a double blind controlled trial. 10th European College of Neuropsychopharmacology Congress. Vienna, Austria. 13th 17th September; 1997. Exclusion Code: X1
347. Trivedi MH, Greer TL, Church TS, et al. Exercise as an augmentation treatment for nonremitted major depressive disorder: A randomized, parallel dose comparison. *J Clin Psychiatry*. 2011;72(5):677-84. Exclusion Code: X5

348. Trivedi MH, Greer TL, Grannemann BD, et al. Exercise as an augmentation strategy for treatment of major depression. *J Psychiatr Pract.* 2006;12(4):205-13. PMID: 2009256065. Language: English. Entry Date: 20070105. Revision Date: 20091218. Publication Type: journal article. Exclusion Code: X5
349. Tsang HWH, Chan EP, Cheung WM. Effects of mindful and non mindful exercises on people with depression: A systematic review. *Br J Clin Psychol.* 2008;47(3):303-22. PMID: 2009-01053-005. First Author & Affiliation: Tsang, Hector W. H. Exclusion Code: X2
350. Turner P, Kantaria R, Young AH. A systematic review and meta-analysis of the evidence base for add-on treatment for patients with major depressive disorder who have not responded to antidepressant treatment: a European perspective. *J Psychopharmacol.* 2014 Feb;28(2):85-98. PMID: 24108407. Exclusion Code: X8
351. van Hees ML, Rotter T, Ellermann T, et al. The effectiveness of individual interpersonal psychotherapy as a treatment for major depressive disorder in adult outpatients: a systematic review. *BMC Psychiatry.* 2013;13:22. PMID: 23312024. Exclusion Code: X8
352. van Schaik A, van Marwijk H, Ader H, et al. Interpersonal psychotherapy for elderly patients in primary care. *Am J Geriatr Psychiatry.* 2006 Sep;14(9):777-86. PMID: 16943174. Exclusion Code: X5
353. Vis PM, Baardewijk M, Einarson TR. Duloxetine and venlafaxine-XR in the treatment of major depressive disorder: a meta-analysis of randomized clinical trials (Structured abstract). *Ann Pharmacother;* 2005. p. 1798-807. Exclusion Code: X2
354. Vitiello B, Shader RI, Parker CB, et al. Hyperforin plasma level as a marker of treatment adherence in the National Institutes of Health Hypericum Depression Trial. *J Clin Psychopharmacol.* 2005 Jun;25(3):243-9. PMID: 15876903. Exclusion Code: X4
355. Vittengl JR, Clark LA, Dunn TW, et al. Reducing relapse and recurrence in unipolar depression: a comparative meta-analysis of cognitive-behavioral therapy's effects. *J Consult Clin Psychol.* 2007 Jun;75(3):475-88. PMID: 17563164. Exclusion Code: X6
356. Wagner JJ. A meta-analysis/literature review comparing the effectiveness of SSRI antidepressants, cognitive behavioral therapy, and placebo for the treatment of depression. US: ProQuest Information & Learning; 2005. Exclusion Code: X6
357. Ward EC. Examining differential treatment effects for depression in racial and ethnic minority women: A qualitative systematic review. *J Natl Med Assoc.* 2007;99(3):265-74. PMID: 2007-03731-005. PMID: 17393951. First Author & Affiliation: Ward, Earlise C. Exclusion Code: X1
358. Warden D, Trivedi MH, Carmody T, et al. Adherence to Antidepressant Combinations and Monotherapy for Major Depressive Disorder: A CO-MED Report of Measurement-Based Care. *J Psychiatr Pract.* 2014 Mar;20(2):118-32. PMID: 24638046. Exclusion Code: X2
359. Watkins ER, Mullan E, Wingrove J, et al. Rumination-focused cognitive-behavioural therapy for residual depression: phase II randomised controlled trial. *Br J Psychiatry.* 2011 Oct;199(4):317-22. PMID: 21778171. Exclusion Code: X5
360. Weisler R, McIntyre RS, Bauer M. Extended-release quetiapine fumarate in the treatment of patients with major depressive disorder: adjunct therapy. *Expert Rev Neurother.* 2013 Nov;13(11):1183-200. PMID: 24175721. Exclusion Code: X1
361. Whiskey E, Werneke U, Taylor D. A systematic review and meta-analysis of *Hypericum perforatum* in depression: A comprehensive clinical review. *Int Clin Psychopharmacol.* 2001;16(5):239-52. PMID: 2001-18595-001. PMID: 11552767. First Author & Affiliation: Whiskey, E.. Release Date: 20010919. Publication Type: Journal, (0100). Exclusion Code: X5

362. Whyte EM, Basinski J, Farhi P, et al. Geriatric depression treatment in nonresponders to selective serotonin reuptake inhibitors. *J Clin Psychiatry*. 2004 Dec;65(12):1634-41. PMID: 15641868. Exclusion Code: X2
363. Wiles N, Thomas L, Abel A, et al. Cognitive behavioural therapy as an adjunct to pharmacotherapy for primary care based patients with treatment resistant depression: Results of the CoBaT randomised controlled trial. *Lancet*. 2013 Feb 2;381(9864):375-84. PMID: 0165506. Exclusion Code: X5
364. Wilkinson P, Alder N, Juszczak E, et al. A pilot randomised controlled trial of a brief cognitive behavioural group intervention to reduce recurrence rates in late life depression. *Int J Geriatr Psychiatry*. 2009 Jan;24(1):68-75. PMID: 18615497. Exclusion Code: X2
365. Williams A-I, Katz D, Ali A, et al. Do essential fatty acids have a role in the treatment of depression? *J Affect Disord*. 2006;93(1-3):117-23. PMID: 2006-07896-014. PMID: 16650900. First Author & Affiliation: Williams, Anna-leila. Exclusion Code: X2
366. Wisniewski SR, Chen CC, Kim E, et al. Global benefit-risk analysis of adjunctive aripiprazole in the treatment of patients with major depressive disorder. *Pharmacoeconom Drug Saf*. 2009 Oct;18(10):965-72. PMID: 19662630. Exclusion Code: X4
367. Wu J, Yeung AS, Schnyer R, et al. Acupuncture for depression: a review of clinical applications. *Can J Psychiatry*. 2012 Jul;57(7):397-405. PMID: 22762294. Exclusion Code: X4
368. Xiao L. A Double-blind, Active-controlled, Randomized Study Comparing Mirtazapine Combined With Paroxetine or Paroxetine Monotherapy in Patients With Major Depressive Patients Without Early Improvement in the First 2 Weeks [NCT01458626]. *Clinicaltrials.gov* [www.clinicaltrials.gov]; 2011. Exclusion Code: X9
369. Xie F, Despiegel N, Danchenko N, et al. Cost effectiveness analysis of escitalopram compared to venlafaxine and fluvoxamine in treatment of major depressive disorder (Structured abstract). *Int J Psychiatry Clin Pract*; 2009. p. 59-69. Exclusion Code: X1
370. Yeung WF, Chung KF, Tso KC, et al. Electroacupuncture for residual insomnia associated with major depressive disorder: a randomized controlled trial. *Sleep*. 2011 Jun;34(6):807-15. PMID: 21629370. Exclusion Code: X2
371. Zajecka J, Dunner DL, Gelenberg AJ, et al. Sexual function and satisfaction in the treatment of chronic major depression with nefazodone, psychotherapy, and their combination. *J Clin Psychiatry*. 2002 Aug;63(8):709-16. PMID: 12197452. Exclusion Code: X2
372. Zhang ZJ, Chen HY, Yip KC, et al. The effectiveness and safety of acupuncture therapy in depressive disorders: systematic review and meta-analysis. *J Affect Disord*. 2010 Jul;124(1-2):9-21. PMID: 19632725. Exclusion Code: X3
373. Zhang ZJ, Ng R, Man SC, et al. Dense cranial electroacupuncture stimulation for major depressive disorder-A single-blind, randomized, controlled study. *PLoS One*. 2012;7(1). Exclusion Code: X3
374. Zilcha-Mano S, Dinger U, McCarthy KS, et al. Changes in well-being and quality of life in a randomized trial comparing dynamic psychotherapy and pharmacotherapy for major depressive disorder. *J Affect Disord*. 2014 Jan;152-154:538-42. PMID: 24176534. Exclusion Code: X3
375. Zu S, Xiang YT, Liu J, et al. A comparison of cognitive-behavioral therapy, antidepressants, their combination and standard treatment for Chinese patients with moderate-severe major depressive disorders. *J Affect Disord*. 2014;152-154(1):262-7. Exclusion Code: X5

Appendix D. Risk of Bias Evaluations

Table D1. Risk of bias domains and ratings

Author, Year Trial Name	Randomi- zation method adequate?	Allocation concealment adequate?	Groups similar at baseline?	Outcome asse- ssors masked?	Care providers masked?	Patients masked?	High overall (i.e., ≥20%) attrition?	High differential (i.e., ≥15%) attrition?	Use Intention- to-treat analyses?	Appropriate method of handling dropouts in analyses used?	Could selective reporting of outcomes be an issue?	Risk of Bias
Barber et al., 2012 ¹	Yes	NR	No	Unclear	No	No	Yes	Yes	Yes	Yes	No	Medium
Bastos et al., 2013 ²	NR	NR	Yes	Yes	No	No	Yes	Yes	Yes	Yes	Yes	Medium
Behnke et al., 2002 ³	Yes	Unclear	Unclear	Unclear	Unclear	Unclear	No	No	Yes	NR	Unclear	Medium
Bjerkénstet et al., 2005 ⁴	Yes	Yes	Yes	Yes	Yes	Yes	No	No	Yes	Yes	No	Low
Blom et al., 2007 ⁵	NR	NR	Yes	Yes	Unclear	Unclear	Yes	No	Unclear	Unclear	No	Medium
Blumenthal et al., 1999 ⁶	Unclear	Unclear	Unclear	Yes	Yes	No	Yes	No	Yes	Yes	No	Medium
Babyak et al., 2000 ⁷												
Blumenthal et al, 2007 ⁸	Yes	Unclear	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Medium
Hoffman et al., 2008 ⁹												
Brenner et al., 2000 ¹⁰	NR	NR	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	High
David et al., 2008 ¹¹	NR	NR	Yes	Yes	No	No	No	No	Yes	Yes	No	Medium
Sava et al., 2009 ¹²												
Davidson et al., 2002 ¹³	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	Unclear	Medium
Dekker et al., 2008 ¹⁴	NR	NR	Yes	NR	No	No	Yes	No	Yes	Yes	No	Medium
DeRubeis et al., 2005 ¹⁵	Yes	Yes	Yes	Yes	No	No	No	No	Yes	Yes	No	Medium
Leykin et al., 2007 ¹⁶												

Table D1. Risk of bias domains and ratings (continued)

Author, Year Trial Name	Randomi- zation method adequate?	Allocation concealment adequate?	Groups similar at baseline?	Outcome asse- ssors masked?	Care providers masked?	Patients masked?	High overall (i.e., ≥20%) attrition?	High differential (i.e., ≥15%) attrition?	Use Intention- to-treat analyses?	Appropriate method of handling dropouts in analyses used?	Could selective reporting of outcomes be an issue?	Risk of Bias
Dimidjian et al., 2006 ¹⁷	NR	NR	Yes	NR	No	No	Yes	Yes	No	NR	No	High
Fava et al., 2005 ¹⁸	NR	NR	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	No	High
Papakostas et al., 2007 ¹⁹												
Frank et al., 2011 ²⁰	NR	NR	No	Unclear	No	No	Yes	No	Yes	Yes	Yes	High
Rucci, 2011 ²¹												
Gastpar et al., 2005 ²²	Yes	Unclear	Yes	Unclear	Yes	Yes	Yes	Yes	Yes	Yes	No	Medium
Gastpar et al., 2006 ²³	Yes	Unclear	Yes	Yes	Yes	No	No	Yes	Yes	No	No	Low
Gertsik et al., 2012 ²⁴	NR	NR	Yes	NR	NR	Yes	Yes	Yes	No	NR	No	High
Harrer et al., 1999 ²⁵	Yes	Unclear	Yes	Unclear	Unclear	Unclear	No	No	No	NR	Unclear	Medium
Hegerl et al., 2010 ²⁶	Yes	Yes	Yes	Yes	No	Yes	No	No	Yes	Yes	Yes	High
Huang et al., 2005 ²⁷	Unclear	Unclear	Unclear	Unclear	No	No	No	No	No	No	No	Medium
Jazayeri et al., 2008 ²⁸	NR	NR	Yes	Yes	Yes	Yes	Yes	No	No	No	No	High
Kennedy et al., 2007 ²⁹	NR	NR	No	NR	No	No	Yes	No	No	NR	No	High
Lam et al., 2013 ³⁰	Yes	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	No	Low
Landenberger et al., 2002 ³¹	NR	NR	No	Yes	No	No	No	No	Yes	Yes	Unclear	Medium
Lenox-Smith and Jiang, 2008 ³²	Yes	Unclear	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	No	Low
McGrath et al., 2013 ³³	Yes	Yes	No	Yes	No	No	Yes	No	No	NR	No	High

Table D1. Risk of bias domains and ratings (continued)

Author, Year Trial Name	Randomi- Zation Method Adequate?	Allocation Concealment Adequate?	Groups Similar at Baseline ?	Outcome Asse- ssors Masked?	Care Providers Masked?	Patients Masked?	High Overall (i.e., ≥20%) Attrition?	High Differential (i.e., ≥15%) Attrition?	Use Intention- to-Treat Analyses?	Appropriate Method of Handling Dropouts in Analyses Used?	Could Selective Reporting of Outcomes be an Issue?	Risk of Bias
Menchetti et al., 2014 ³⁴	Yes	Yes	Yes	Yes	No	No	No	No	Yes	Yes	Yes	Medium
Miranda et al., 2003 ³⁵	Yes	Yes	Yes	Yes	No	No	Unclear	No	Yes	Yes	No	Medium
WECare												
Mischoulon et al., 2014 ³⁶	Yes	NR	NR	Yes	Yes	Yes	Yes	No	Yes	Yes	No	High
Moradveisi et al., 2013 ³⁷	Yes	NR	Yes	Yes	No	No	Yes	Yes	Yes	Yes	No	High
Moreno et al., 2006 ³⁸	Unclear	Unclear	No	Unclear	Yes	Yes	No	No	No	Yes	Yes	High
Mynors-Wallis et al., 2000 ³⁹	Unclear	Unclear	Yes	Yes	No	No	Yes	Yes	Unclear	Unclear	Yes	Medium
Qu et al., 2013 ⁴⁰	Yes	Unclear	Yes	Yes	No	No	No	No	No	NR	No	Medium
Chen et al., 2014 ⁴¹												
Raue et al., 2009 ⁴²	NR	NR	NR	NR	No	No	No	NR	NR	NR	Yes	High
Salminen et al., 2008 ⁴³	NR	NR	Yes	NR	No	No	Yes	No	Yes	NR	No	Medium
Kronstrom et al., 2009 ⁴⁴												
Schrader et al., 2000 ⁴⁵	Unclear	Unclear	Unclear	Unclear	Yes	Yes	No	No	Yes	Yes	No	Medium
Segal et al., 2006 ⁴⁶	NR	NR	Yes	NR	No	No	Yes	Yes	No	NR	No	High
Shamsaei et al., 2008 ⁴⁷	Yes	NR	Yes	No	No	No	NR	NR	NR	NR	No	High
Song et al., 2007 ⁴⁸	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	NR	Unclear	Yes	High
Sun et al., 2013 ⁴⁹	Yes	Yes	Unclear	Unclear	No	Yes	No	No	Yes	No	No	High

Table D1. Risk of bias domains and ratings (continued)

Author, Year Trial Name	Randomi- Zation Method Adequate?	Allocation Concealment Adequate?	Groups Similar at Baseline ?	Outcome Asse- ssors Masked?	Care Providers Masked?	Patients Masked?	High Overall (i.e., ≥20%) Attrition?	High Differential (i.e., ≥15%) Attrition?	Use Intention- to-Treat Analyses?	Appropriate Method of Handling Dropouts in Analyses Used?	Could Selective Reporting of Outcomes be an Issue?	Risk of Bias
Szegedi et al., 2005 ⁵⁰	Yes	Yes	Yes	NR	Yes	Yes	No	Yes	Yes	No	No	Medium
Thase et al., 2007 ⁵¹	Yes	Yes	Yes	Yes	No	No	Yes	Unclear	Yes	Yes	No	Medium
Rush et al., 2006 ⁵²												
Trivedi et al., 2006 ⁵³												
STAR*D												
van Gurp et al., 2002 ⁵⁴	Yes	NR	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	No	Medium
Zhang et al., 2009 ⁵⁵	Yes	Yes	Yes	Yes	No	Yes	No	No	Yes	Yes	No	Medium

Appendix D References

1. Barber JP, Barrett MS, Gallop R, et al. Short-term dynamic psychotherapy versus pharmacotherapy for major depressive disorder: a randomized, placebo-controlled trial. *J Clin Psychiatry*. 2012 Jan;73(1):66-73. Epub: 2011/12/14. PMID: 22152401.
2. Bastos AG, Guimaraes LS, Trentini CM. Neurocognitive changes in depressed patients in psychodynamic psychotherapy, therapy with fluoxetine and combination therapy. *J Affect Disord*. 2013 Dec;151(3):1066-75. PMID: 24103853.
3. Behnke K, Jensen GS, Graubaum HJ, et al. Hypericum perforatum versus fluoxetine in the treatment of mild to moderate depression. *Adv Ther*. 2002 Jan-Feb;19(1):43-52. Epub: 2002/05/15. PMID: 12008860.
4. Bjerkenstedt L, Edman GV, Alken RG, et al. Hypericum extract LI 160 and fluoxetine in mild to moderate depression: a randomized, placebo-controlled multi-center study in outpatients. *Eur Arch Psychiatry Clin Neurosci*. 2005 Feb;255(1):40-7. Epub: 2004/11/13. PMID: 15538592.
5. Blom MB, Jonker K, Dusseldorp E, et al. Combination treatment for acute depression is superior only when psychotherapy is added to medication. *Psychother Psychosom*. 2007;76(5):289-97. Epub: 2007/08/19. PMID: 17700049.
6. Blumenthal JA, Babyak MA, Moore KA, et al. Effects of exercise training on older patients with major depression. *Arch Intern Med*. 1999 Oct 25;159(19):2349-56. Epub: 1999/11/05. PMID: 10547175.
7. Babyak M, Blumenthal JA, Herman S, et al. Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. *Psychosom Med*. 2000 Sep-Oct;62(5):633-8. Epub: 2000/10/06. PMID: 11020092.
8. Blumenthal JA, Babyak MA, Doraiswamy PM, et al. Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med*. 2007 Sep-Oct;69(7):587-96. Epub: 2007/09/12. PMID: 17846259.
9. Hoffman BM, Blumenthal JA, Babyak MA, et al. Exercise fails to improve neurocognition in depressed middle-aged and older adults. *Med Sci Sports Exerc*. 2008 Jul;40(7):1344-52. Epub: 2008/06/27. PMID: 18580416.
10. Brenner R, Azbel V, Madhusoodanan S, et al. Comparison of an extract of hypericum (LI 160) and sertraline in the treatment of depression: a double-blind, randomized pilot study. *Clin Ther*. 2000 Apr;22(4):411-9. PMID: 10823363.
11. David D, Szentagotai A, Lupu V, et al. Rational emotive behavior therapy, cognitive therapy, and medication in the treatment of major depressive disorder: a randomized clinical trial, posttreatment outcomes, and six-month follow-up. *J Clin Psychol*. 2008 Jun;64(6):728-46. Epub: 2008/05/14. PMID: 18473339.
12. Sava FA, Yates BT, Lupu V, et al. Cost-effectiveness and cost-utility of cognitive therapy, rational emotive behavioral therapy, and fluoxetine (Prozac) in treating depression: a randomized clinical trial. *J Clin Psychol*. 2009 Jan;65(1):36-52. Epub: 2008/12/04. PMID: 19051275.
13. Davidson JRT, Gadde KM, Fairbank JA, et al. Effect of *Hypericum perforatum* (St John's wort) in major depressive disorder: a randomized controlled trial. *J Am Med Assoc*. 2002;287(14):1807-14.
14. Dekker JJ, Koelen JA, Van HL, et al. Speed of action: the relative efficacy of short psychodynamic supportive psychotherapy and pharmacotherapy in the first 8 weeks of a treatment algorithm for depression. *J Affect Disord*. 2008 Jul;109(1-2):183-8. Epub: 2007/12/07. PMID: 18061276.
15. DeRubeis RJ, Hollon SD, Amsterdam JD, et al. Cognitive therapy vs medications in the treatment of moderate to severe depression. *Arch Gen Psychiatry*. 2005 Apr;62(4):409-16. Epub: 2005/04/06. PMID: 15809408.

16. Leykin Y, Amsterdam JD, DeRubeis RJ, et al. Progressive resistance to a selective serotonin reuptake inhibitor but not to cognitive therapy in the treatment of major depression. *J Consult Clin Psychol*. 2007 Apr;75(2):267-76. Epub: 2007/05/02. PMID: 17469884.
17. Dimidjian S, Hollon SD, Dobson KS, et al. Randomized trial of behavioral activation, cognitive therapy, and antidepressant medication in the acute treatment of adults with major depression. *J Consult Clin Psychol*. 2006 Aug;74(4):658-70. Epub: 2006/08/03. PMID: 16881773.
18. Fava M, Alpert J, Nierenberg AA, et al. A double-blind, randomized trial of St John's wort, fluoxetine, and placebo in major depressive disorder. *J Clin Psychopharmacol*. 2005 Oct;25(5):441-7. Epub: 2005/09/15. PMID: 16160619.
19. Papakostas GI, Crawford CM, Scalia MJ, et al. Timing of clinical improvement and symptom resolution in the treatment of major depressive disorder. A replication of findings with the use of a double-blind, placebo-controlled trial of Hypericum perforatum versus fluoxetine. *Neuropsychobiology*. 2007;56(2-3):132-7. Epub: 2008/02/09. PMID: 18259086.
20. Frank E, Cassano GB, Rucci P, et al. Predictors and moderators of time to remission of major depression with interpersonal psychotherapy and SSRI pharmacotherapy. *Psychol Med*. 2011 Jan;41(1):151-62. Epub: 2010/04/13. PMID: 20380782.
21. Rucci P, Frank E, Scocco P, et al. Treatment-emergent suicidal ideation during 4 months of acute management of unipolar major depression with SSRI pharmacotherapy or interpersonal psychotherapy in a randomized clinical trial. *Depress Anxiety*. 2011 Apr;28(4):303-9. Epub: 2011/02/11. PMID: 21308882.
22. Gastpar M, Singer A, Zeller K. Efficacy and tolerability of hypericum extract STW3 in long-term treatment with a once-daily dosage in comparison with sertraline. *Pharmacopsychiatry*. 2005 Mar;38(2):78-86. PMID: 15744631.
23. Gastpar M, Singer A, Zeller K. Comparative efficacy and safety of a once-daily dosage of hypericum extract STW3-VI and citalopram in patients with moderate depression: a double-blind, randomised, multicentre, placebo-controlled study. *Pharmacopsychiatry*. 2006 Mar;39(2):66-75. Epub: 2006/03/24. PMID: 16555167.
24. Gertsik L, Poland RE, Bresee C, et al. Omega-3 fatty acid augmentation of citalopram treatment for patients with major depressive disorder. *J Clin Psychopharmacol*. 2012 Feb;32(1):61-4. Epub: 2011/12/27. PMID: 22198441.
25. Harrer G, Schmidt U, Kuhn U, et al. Comparison of equivalence between the St. John's wort extract LoHyp-57 and fluoxetine. *Arzneimittelforschung*. 1999 Apr;49(4):289-96. Epub: 1999/05/25. PMID: 10337446.
26. Hegerl U, Hautzinger M, Mergl R, et al. Effects of pharmacotherapy and psychotherapy in depressed primary-care patients: a randomized, controlled trial including a patients' choice arm. *Int J Neuropsychopharmacol*. 2010;13(1):31-44. PMID: 19341510.
27. Huang Y, Htut W, Li D, et al. Studies on the clinical observation and cerebral glucose metabolism in depression treated by electro-scalp acupuncture compared to fluoxetine. *Int J Clin Acupunct*. 2005;14(1):7-26. PMID: 0077160.
28. Jazayeri S, Tehrani-Doost M, Keshavarz SA, et al. Comparison of therapeutic effects of omega-3 fatty acid eicosapentaenoic acid and fluoxetine, separately and in combination, in major depressive disorder. *Aust N Z J Psychiatry*. 2008 Mar;42(3):192-8. Epub: 2008/02/06. PMID: 18247193.
29. Kennedy SH, Konarski JZ, Segal ZV, et al. Differences in brain glucose metabolism between responders to CBT and venlafaxine in a 16-week randomized controlled trial. *Am J Psychiatry*. 2007 May;164(5):778-88. Epub: 2007/05/04. PMID: 17475737.
30. Lam RW, Parikh SV, Ramasubbu R, et al. Effects of combined pharmacotherapy and psychotherapy for improving work functioning in major depressive disorder. *Br J Psychiatry*. 2013 Nov;203(5):358-65. PMID: 24029535.

31. Landenberger NAD. Self-concept and attributional style in the treatment of depression: ProQuest Information & Learning; 2002.
32. Lenox-Smith AJ, Jiang Q. Venlafaxine extended release versus citalopram in patients with depression unresponsive to a selective serotonin reuptake inhibitor. *Int Clin Psychopharmacol*. 2008 May;23(3):113-9. Epub: 2008/04/15. PMID: 18408525.
33. McGrath CL, Kelley ME, Holtzheimer PE, et al. Toward a neuroimaging treatment selection biomarker for major depressive disorder. *JAMA Psychiatry*. 2013 Aug;70(8):821-9. Epub: 2013/06/14. PMID: 23760393.
34. Menchetti M, Rucci P, Bortolotti B, et al. Moderators of remission with interpersonal counselling or drug treatment in primary care patients with depression: randomised controlled trial. *Br J Psychiatry*. 2014 Feb;204(2):144-50. PMID: 24311553.
35. Miranda J, Chung JY, Green BL, et al. Treating depression in predominantly low-income young minority women: a randomized controlled trial. *JAMA*. 2003 Jul 2;290(1):57-65. Epub: 2003/07/03. PMID: 12837712.
36. Mischoulon D, Price LH, Carpenter LL, et al. A double-blind, randomized, placebo-controlled clinical trial of S-adenosyl-L-methionine (SAME) versus escitalopram in major depressive disorder. *J Clin Psychiatry*. 2014 Dec 24. Epub: 2014/02/07. PMID: 24500245.
37. Moradveisi L, Huibers MJ, Renner F, et al. Behavioural activation v. antidepressant medication for treating depression in Iran: randomised trial. *Br J Psychiatry*. 2013 Mar;202(3):204-11. Epub: 2013/02/09. PMID: 23391727.
38. Moreno RA, Teng CT, Almeida KM, et al. *Hypericum perforatum* versus fluoxetine in the treatment of mild to moderate depression: a randomized double-blind trial in a Brazilian sample. *Rev Bras Psiquiatr*. 2006 Mar;28(1):29-32. PMID: 16612487.
39. Mynors-Wallis LM, Gath DH, Day A, et al. Randomised controlled trial of problem solving treatment, antidepressant medication, and combined treatment for major depression in primary care. *BMJ*. 2000 Jan 1;320(7226):26-30. Epub: 2000/01/05. PMID: 10617523.
40. Qu SS, Huang Y, Zhang ZJ, et al. A 6-week randomized controlled trial with 4-week follow-up of acupuncture combined with paroxetine in patients with major depressive disorder. *J Psychiatr Res*. 2013 Jun;47(6):726-32. Epub: 2013/03/19. PMID: 23498306.
41. Chen JQ, Lin WR, Wang SX, et al. Acupuncture/electroacupuncture enhances antidepressant effect of seroxat: the symptom checklist-90 scores. *Neural Regen Res*. 2014;9(2):213-22.
42. Raue PJ, Schulberg HC, Heo M, et al. Patients' depression treatment preferences and initiation, adherence, and outcome: a randomized primary care study. *Psychiatr Serv*. 2009 Mar;60(3):337-43. Epub: 2009/03/03. PMID: 19252046.
43. Salminen JK, Karlsson H, Hietala J, et al. Short-term psychodynamic psychotherapy and fluoxetine in major depressive disorder: a randomized comparative study. *Psychother Psychosom*. 2008;77(6):351-7. Epub: 2008/08/15. PMID: 18701831.
44. Kronstrom K, Salminen JK, Hietala J, et al. Does defense style or psychological mindedness predict treatment response in major depression? *Depress Anxiety*. 2009;26(7):689-95. Epub: 2009/06/06. PMID: 19496102.
45. Schrader E. Equivalence of St John's wort extract (Ze 117) and fluoxetine: a randomized, controlled study in mild-moderate depression. *Int Clin Psychopharmacol*. 2000 Mar;15(2):61-8. PMID: 10759336.
46. Segal ZV, Kennedy S, Gemar M, et al. Cognitive reactivity to sad mood provocation and the prediction of depressive relapse. *Arch Gen Psychiatry*. 2006 Jul;63(7):749-55. Epub: 2006/07/05. PMID: 16818864.

47. Shamsaei F, Rahimi A, Zarabian MK, et al. Efficacy of pharmacotherapy and cognitive therapy, alone and in combination in major depressive disorder. *Hong Kong J Psychiatr*. 2008;18(2):76-80.
48. Song Y, Zhou D, Fan J, et al. Effects of electroacupuncture and fluoxetine on the density of GTP-binding-proteins in platelet membrane in patients with major depressive disorder. *J Affect Disord*. 2007 Mar;98(3):253-7. Epub: 2006/08/22. PMID: 16919758.
49. Sun H, Zhao H, Ma C, et al. Effects of electroacupuncture on depression and the production of glial cell line-derived neurotrophic factor compared with fluoxetine: a randomized controlled pilot study. *J Altern Complement Med*. 2013 Sep;19(9):733-9. PMID: 23647408.
50. Szegedi A, Kohnen R, Dienel A, et al. Acute treatment of moderate to severe depression with hypericum extract WS 5570 (St John's wort): randomised controlled double blind non-inferiority trial versus paroxetine. *BMJ*. 2005 Mar 5;330(7490):503. Epub: 2005/02/15. PMID: 15708844.
51. Thase ME, Friedman ES, Biggs MM, et al. Cognitive therapy versus medication in augmentation and switch strategies as second-step treatments: a STAR*D report. *Am J Psychiatry*. 2007 May;164(5):739-52. Epub: 2007/05/04. PMID: 17475733.
52. Rush AJ, Trivedi MH, Wisniewski SR, et al. Bupropion-SR, sertraline, or venlafaxine-XR after failure of SSRIs for depression. *N Engl J Med*. 2006 Mar 23;354(12):1231-42. Epub: 2006/03/24. PMID: 16554525.
53. Trivedi MH, Fava M, Wisniewski SR, et al. Medication augmentation after the failure of SSRIs for depression. *N Engl J Med*. 2006 Mar 23;354(12):1243-52. Epub: 2006/03/24. PMID: 16554526.
54. van Gurp G, Meterissian GB, Haiek LN, et al. St John's wort or sertraline? Randomized controlled trial in primary care. *Can Fam Physician*. 2002 May;48:905-12. Epub: 2002/06/11. PMID: 12053635.
55. Zhang WJ, Yang XB, Zhong BL. Combination of acupuncture and fluoxetine for depression: a randomized, double-blind, sham-controlled trial. *J Altern Complement Med*. 2009 Aug;15(8):837-44. Epub: 2009/08/15. PMID: 19678773.

Appendix E: Summary of Findings Tables

KQ 1 and KQ4: Summary of Findings Tables

Table 1. Benefits of second-generation antidepressants compared with cognitive behavioral therapy monotherapy

Outcomes	Anticipated absolute effects ^a : <i>Benefit with CBT monotherapy</i>	Anticipated absolute effects ^a : (95% CI): <i>Benefit with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Response Assessed with: HAM-D Followup: range 8 to 16 weeks	45 per 100	47 per 100 (40 to 55)	RR, 1.03 (0.87 to 1.21)	607 (5 trials)	Moderate ^b	Comparisons limited to fluoxetine, fluvoxamine, paroxetine, or sertraline and CBT, CT, PST, or REBT. Sensitivity analysis with 3 additional trials (rated high risk of bias) did not change the statistical significance of the results (RR, 1.09; 95% CI, 0.99 to 1.20).
Remission Assessed with: HAM-D Followup: range 12 to 16 weeks	49 per 100	59 per 100 (48 to 72)	RR, 1.19 (0.98 to 1.45)	379 (3 trials)	Low ^{b, c}	Comparisons limited to fluoxetine, fluvoxamine, or paroxetine and CBT, CT, PST, or REBT sensitivity analysis with 3 additional trials (rated high risk of bias) resulted in a statistically significant difference favoring SGA (RR, 1.19; 95% CI, 1.04 to 1.33).
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity Assessed with: Social Adjustment Scale Follow up: mean 12 weeks	Mean scores were within 0.3 points between groups and CIs overlapped.	Mean scores were within 0.3 points between groups and CIs overlapped.	Not estimable	116 (1 study)	Low ^{d, e}	Comparison limited to fluvoxamine or paroxetine and PST.

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for inconsistency: inconsistent direction of point estimates.

^c Downgraded for suspected selective outcome reporting bias.

^d Downgraded for imprecision: sample size that does not fulfill optimal information size (OIS).

^e Downgraded for risk of bias: outcomes reporting bias; most trials did not report on functional capacity.

CBT = cognitive behavioral therapy; CI = confidence interval; CT = cognitive therapy; HAM-D = Hamilton Depression Rating Scale; NR: not reported; PST = problem solving therapy; REBT = rational emotive behavior therapy; RR = risk ratio; SGA = second-generation antidepressant

Table 2. Benefits of second-generation antidepressants compared with combinations of second-generation antidepressants and cognitive behavioral therapy

Outcomes	Anticipated absolute effects ^a : <i>Benefit with combination of SGA and CBT</i>	Anticipated absolute effects ^a : (95% CI): <i>Benefit with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of Evidence	Comments
Response Assessed with: MADRS or HAM-D Follow up: mean 12 weeks	68 per 100	70 per 100 (58 to 85)	RR, 1.03 (0.85 to 1.26)	174 (2 trials)	Low ^{b, c}	Comparison limited to escitalopram, fluvoxamine, or paroxetine and problem solving therapy or telephone CBT.
Remission Assessed with: MADRS or HAM-D Follow up: mean 12 weeks	55 per 100	58 per 100 (45 to 76)	RR, 1.06 (0.82 to 1.38)	174 (2 trials)	Low ^{b, c}	Comparison limited to escitalopram, fluvoxamine, or paroxetine and problem solving therapy or telephone CBT.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity Assessed with: Multiple scales Follow up: mean 12 weeks	Patients receiving the combination reported greater improvement on 3 of 5 work functioning measures compared with patients on SGA alone	Patients receiving the combination reported greater improvement on 3 of 5 work functioning measures compared with patients on SGA alone	Not estimable	170 (2 trials)	Low ^{b, c}	Comparison limited to escitalopram, fluvoxamine, or paroxetine and problem solving therapy or telephone CBT.

^a The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for inconsistency: inconsistent direction of point estimates.

^c Downgraded for imprecision: sample size that does not fulfill optimal information size (OIS).

CBT = cognitive behavioral therapy; CI = confidence interval; MADRS = Montgomery-Åsberg Depression Rating Scale; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 3. Benefits of second-generation antidepressants compared with integrative therapies monotherapy

Outcomes	Anticipated absolute effects ^a : <i>Benefit with integrative therapies alone</i>	Anticipated absolute effects ^a : (95% CI): <i>Benefit with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of Evidence	Comments
Response Assessed with: HAM-D Follow up: mean 6 weeks	61 per 100	62 per 100 (53 to 75)	RR, 1.02 (0.86 to 1.22)	318 (1 trial)	Low ^{b, c}	Comparison limited to escitalopram and IPT.
Remission Assessed with: HAM-D Followup: range 8 to 12 weeks	50 per 100	46 per 100 (39 to 54)	RR, 0.92 (0.78 to 1.08)	605 (2 trials)	Low ^{d, e}	Comparison limited to citalopram, escitalopram, or sertraline and IPT. A third study (rated high risk of bias) reported no significant difference in effect but did not present rates of remission.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^aThe risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: sample size that does not fulfill optimal information size (OIS).

^c Downgraded for risk of bias: high risk of bias due to insufficient reporting of methods and baseline differences between groups in duration of illness.

^d Downgraded for inconsistency: inconsistent direction of point estimates.

^e Downgraded for risk of bias: one of the trials was rated high risk of bias due to insufficient reporting of methods and baseline differences between groups in duration of illness.

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; IPT = Interpersonal Psychotherapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 4. Benefits of second-generation antidepressants compared with combinations of second-generation antidepressants and integrative therapies

Outcomes	Anticipated absolute effects ^a : <i>Benefit with combination of SGA and integrative therapies</i>	Anticipated absolute effects ^a : (95% CI): <i>Benefit with SGA</i>	Relative effect (95% CI) ^a	Number of participants (Trials)	Strength of Evidence	Comments
Response	NA	NA	NA	0 (0 trials)	Insufficient	None
Remission Assessed with: HAM-D Followup: range 8 to 12 weeks	NR	NR	OR, 3.22 (1.02 to 10.12)	97 (1 trial)	Low ^{b, c}	Comparison limited to nefazodone and combination of nefazodone and IPT.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: sample size that does not fulfill optimal information size (OIS).

^c Further downgraded for imprecision (very wide confidence interval).

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; IPT = NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 5. Benefits of second-generation antidepressants compared with psychodynamic therapies monotherapy

Outcomes	Anticipated absolute effects ^a : <i>Benefit or psychodynamic therapies alone</i>	Anticipated absolute effects ^a : (95% CI): <i>Benefit or risk with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of Evidence	Comments
Response	NA	NA	NA	0 (0 trials)	Insufficient	None
Remission assessed with: HAM-D followup: mean 16 weeks	46 per 100	48 per 100 (27 to 86)	RR, 1.04 (0.58 to 1.86)	51 (1 trial)	Low ^{b, c}	Comparison limited to fluoxetine.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	Few statistically significant differences in various scales. In one study, the proportion of patients on sick leave was higher in the SGA group than the PSYD group (12% vs. 4%, not significant),	Few statistically significant differences in various scales. In one study, the proportion of patients on sick leave was higher in the SGA group than the PSYD group (12% vs. 4%, not significant)	Not estimable	221 (2 trials)	Low ^{b, d}	Comparison limited to fluoxetine.

^a The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: sample size that does not fulfill optimal information size (OIS).

^c Downgraded for risk of bias: outcomes reporting bias; most trials did not report on remission

^d Downgraded for inconsistency: inconsistent direction of point estimates.

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; NR = not reported; PSYD = psychodynamic therapy; RR = Risk ratio; SGA = second-generation antidepressant

Table 6. Benefits of second-generation antidepressants compared with combinations of second-generation antidepressants and psychodynamic therapies

Outcomes	Anticipated absolute effects ^a : <i>Benefit with integrative therapies alone</i>	Anticipated absolute effects ^a : (95% CI): <i>Benefit with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of Evidence	Comments
Response	NA	NA	NA	0 (0 trials)	Insufficient	None
Remission	NA	NA	NA	0 (0 trials)	Insufficient	None
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	Effects on WAIS-III measures were similar for SGA and the combination of SGA and PSYD.	Effects on WAIS-III measures were similar for SGA and the combination of SGA and PSYD.	Not estimable	272 (1 trial)	Low ^{b, c}	None

^aThe risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision (single study, small sample size, unable to estimate an overall effect).

^c Further downgraded for imprecision (wide range of effects for various measures).

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; IPT = NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 7. Benefits of second-generation antidepressants compared with third-wave cognitive behavioral therapy monotherapy

Outcomes	Anticipated absolute effects ^a : <i>Benefit with combination of SGA and third wave CBT</i>	Anticipated absolute effects ^a : (95% CI): <i>Benefit with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of Evidence	Comments
Response Assessed with: HAM-D Follow up: mean 13 weeks	88 per 100	66 per 100 (53 to 83)	RR, 0.75 (0.6 to 0.94)	100 (1 trial)	Insufficient ^{b, c, d}	Comparison limited to sertraline.
Remission Assessed with: HAM-D Follow up: mean 13 weeks	82 per 100	48 per 100 (34 to 66)	RR, 0.59 (0.42 to 0.8)	100 (1 trial)	Insufficient ^{b, c, d}	Comparison limited to sertraline.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^aThe risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: sample size that does not fulfill optimal information size (OIS).

^c Downgraded for risk of bias: high overall and differential attrition not accounted for in reported effect size.

^d Downgraded for risk of bias: suspected reporting bias.

CBT = cognitive behavioral therapy; CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; NR: not reported; RR: Risk ratio; SGA = second-generation antidepressant

Table 8. Benefits of second-generation antidepressants compared with acupuncture monotherapy

Outcomes	Anticipated Absolute Effects^a: <i>Benefit with Acupuncture</i>	Anticipated Absolute Effects^a: (95% CI): <i>Benefit with SGA</i>	Relative Effect (95% CI)	Number of Participants (Trials)	Strength of Evidence	Comments
Response Assessed with: HAM-D followup: mean 6 weeks	64 per 100	62 per 100 (50 to 77)	RR, 0.97 (0.78 to 1.21)	184 (2 trials)	Low ^{b, c}	Direct evidence limited to comparisons of fluoxetine vs acupuncture. Results consistent with NWMA comparisons to SGA medications (RR, 0.80, 95% CI, 0.46-1.40).
Remission Assessed with: HAM-D followup: mean 6 weeks	NA	NA	NA	0 (0 trials)	Insufficient	None
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for high risk of bias: high dropout; uncertainty about randomization and allocation concealment; no masking of outcome assessors

^c Downgraded for imprecision: few events not meeting optimal information size (OIS)

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; NWMA = network meta-analysis; RR: Risk ratio; SGA = second-generation antidepressant

Table 9. Benefits of second-generation antidepressants compared with combination of SGA and acupuncture

Outcomes	Anticipated Absolute Effects^a: <i>Benefit with Combination of SGA and Acupuncture</i>	Anticipated Absolute Effects^a: (95% CI): <i>Benefit with SGA</i>	Relative Effect (95% CI)	Number of Participants (Trials)	Strength of Evidence	Comments
Response Assessed with: HAM-D followup: mean 6 weeks	71 per 100	52 per 100 (43 to 63)	RR, 0.73 (0.61 to 0.89)	288 (2 trials)	Low ^{b, c}	Direct evidence limited to two studies comparing either fluoxetine or paroxetine with acupuncture plus SGA.
Remission Assessed with: HAM-D followup: mean 6 weeks	28 per 100	23 per 100 (12 to 42)	RR, 0.83 (0.45 to 1.53)	157 (1 trial)	Low ^b	Direct evidence limited to a single trial of paroxetine with combined acupuncture plus paroxetine.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^aThe risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: few events

^c Downgraded for inconsistency: large effect size differences between studies

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; RR: Risk ratio; SGA = second-generation antidepressant

Table 10. Benefits of second-generation antidepressants compared with omega-3 fatty acids monotherapy

Outcomes	Anticipated Absolute Effects^a: Benefit with Omega-3 Fatty Acids	Anticipated Absolute Effects^a (95% CI): Benefit with SGA	Relative Effect (95% CI)	Number of Participants (Trials)	Strength of Evidence	Comments
Response Assessed with: HAM-D Follow up: mean 8 weeks	45 per 100	40 per 100 (19 to 82)	RR, 0.5 (0.24 to 1.04)	NA, Network meta-analysis	Low ^{a,b}	Direct evidence is limited to a comparison of fluoxetine with Omega-3 fatty acids. Results from network meta-analyses conflict with findings of the RCT and indicate greater efficacy of SGAs (RR, 2.00; 95% CI, 0.96 to 4.18) compared with Omega-3 fatty acids
Remission Assessed with: HAM-D Follow up: mean 8 weeks	NA	NA	NA	0 (0 trials)	Insufficient	None
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for indirectness: results are based on network meta-analyses

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; RR: Risk ratio; SGA = second-generation antidepressant

Table 11. Benefits of second-generation antidepressants compared with combination of SGA and omega-3 fatty acids

Outcomes	Anticipated Absolute Effects^a: <i>Benefit with combination of SGA and Omega-3 Fatty Acids</i>	Anticipated Absolute Effects^a: (95% CI): <i>Benefit with SGA</i>	Relative Effect (95% CI)	Number of Participants (Trials)	Strength of Evidence	Comments
Response Assessed with: HAM-D Follow up mean 8 weeks	47 per 100	29 per 100 (16 to 54)	RR, 0.62 (0.33 to 1.14)	72 (2 trials)	Insufficient ^{b, c}	Direct evidence is limited to a comparison of fluoxetine with omega-3 fatty acids..
Remission Assessed with: HAM-D Follow up mean 8 weeks	44 per 100	18 per 100 (7 to 51)	RR, 0.41 (0.15 to 1.14)	42 (1 trial)	Insufficient ^{b, c}	Direct evidence is limited to a comparison of fluoxetine with omega-3 fatty acids.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for high risk of bias: high attrition and lack of ITT analysis

^c Downgraded for imprecision: few events and confidence interval crosses threshold of appreciable difference

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; RR: Risk ratio; SGA = second-generation antidepressant

Table 12. Benefits of second-generation antidepressants compared with S-Adenosyl methionine monotherapy

Outcomes	Anticipated Absolute Effects^a: <i>Benefit with combination of SGA and Integrative Therapy</i>	Anticipated Absolute Effects^a: (95% CI): <i>Benefit with SGA</i>	Relative Effect (95% CI)	Number of Participants (Trials)	Strength of Evidence	Comments
Response Assessed with: HAM-D Follow up mean 12 weeks	36 per 100	34 per 100 (21 to 54)	RR, 1.22 (0.66 to 2.27)	NA; results based on network meta-analyses	Low ^{b, c}	Direct evidence is limited to a single comparison of escitalopram with SAME. Network meta-analyses found no statistically significant differences in response rates between SGA estimate: 1.22 (0.66, 2.27) compared with SAME.
Remission Assessed with: HAM-D Follow up mean 12 weeks	28 per 100	28 per 100 (16 to 48)	RR, 0.98 (0.57 to 1.71)	129 (1 trial)	Insufficient ^{b, c}	Direct evidence is limited to a single comparison of escitalopram with SAME.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: small study size

^c Downgraded for indirectness: results are based on network meta-analyses

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; NR: not reported; RR: Risk ratio; SAME = S-Adenosylmethionine; SGA = second-generation antidepressant

Table 13. Benefits of second-generation antidepressants compared with St. John's wort monotherapy

Outcomes	Anticipated Absolute Effects^a: Benefit with St. John's Wort	Anticipated Absolute Effects^a (95% CI): Benefit with SGA	Relative Effect (95% CI)	Number of Participants (Trials)	Strength of Evidence	Comments
Response Assessed with: HAM-D Follow up: range 4-12 weeks	57 per 100	55 per 100 (47 to 65)	RR, 0.97 (0.82 to 1.15)	1430 (8 trials)	Moderate ^b	Evidence is based on the comparison of SSRIs with SJW. Results from NWMA demonstrate reduced response rate with SGAs compared to SJW (RR, 0.82; 95% CI, 0.69 to 0.98).
Response in subgroup of older adults Assessed with: HAM-D Follow up: mean 6 weeks	65 per 100	68 per 100 (55 to 84)	RR, 1.05 (0.84 to 1.3)	161 (1 trial)	Low ^c	Comparison limited to fluoxetine and SJW in older adults (60 to 80 years)
Remission Assessed with: HAM-D follow up: mean 13 weeks	25 per 100	21 per 100 (17 to 26)	RR, 0.83 (0.66 to 1.04)	683 (4 trials)	Moderate ^d	Evidence is based on the comparison of SSRIs with SJW.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^aThe risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for inconsistency: moderate heterogeneity (I-squared 56 percent)

^cDowngraded for very serious imprecision

^d Downgraded for imprecision: few events overall and confidence interval crosses threshold of appreciable difference

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; NR: not reported; NWMA = network meta-analyses; RR: Risk ratio; SGA = second-generation antidepressant; SJW = St. John's wort; SSRI = selective serotonin reuptake inhibitor

Table 14. Benefits of second-generation antidepressants compared with exercise monotherapy

Outcomes	Anticipated absolute effects^a: <i>Benefit with exercise</i>	Anticipated absolute effects^a: (95% CI) : <i>Benefit with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of Evidence	Comments
Response Assessed with: HAM-D17 Follow up: mean 16 weeks	82 per 100	94 per 100 (41 to 217)	RR, 1.87 (0.81 to 4.33) ^d	Not estimable (network meta-analyses of RCTs)	Low ^{b, c}	Estimates based on network meta-analyses.
Remission Assessed with: HAM-D17 <8, no longer meeting criteria for MDD Follow up: mean 16 weeks	50 per 100	55 per 100 (44 to 70)	RR, 1.1 (0.87 to 1.39) ^d	254 (2 trials)	Moderate ^b	Comparison is limited to sertraline versus exercise.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: few events, confidence intervals cross threshold of appreciable difference

^c Downgraded for indirectness: estimates are based on network meta-analyses

^d Crude RR

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; MDD = major depressive disorder; NR = not reported; RR: Risk ratio; SGA = second-generation antidepressant

Table 15. Benefits of second-generation antidepressants compared with combination of second-generation antidepressants and exercise

Outcomes	Anticipated absolute effects^a: <i>Benefit with combination of SGA and exercise</i>	Anticipated absolute effects^a (95% CI) : <i>Benefit with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of Evidence	Comments
Response	NR	NR	Not estimable	0 (0 trial)	Insufficient	None
Remission Assessed with: HAM-D17 and no longer meeting criteria for MDD Follow up: mean 16 weeks	66 per 100	69 per 100 (52 to 90)	RR, 1.05 (0.8 to 1.03) ^c	103 (1 trial)	Low ^b	Comparison is limited to sertraline versus sertraline plus exercise.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: few events, confidence intervals cross threshold of appreciable difference

^c Crude RR

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; MDD = major depressive disorder; NR = not reported; RR: Risk ratio; SGA = second-generation antidepressant

Table 16. Benefits of second-generation antidepressants compared with cognitive behavioral therapy as a function of severity

Outcomes	Anticipated absolute effects^a: <i>Benefit with CBT monotherapy</i>	Anticipated absolute effects^a: (95% CI): <i>Benefit with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Response to treatment for high severity patients Assessed with: HAM-D Follow up: mean 16 weeks	56 per 100	40 per 100 (25 to 64)	RR, 0.72 (0.45 to 1.15)	82 (1 trial)	Insufficient ^{b, c}	Comparisons limited to paroxetine and CT.
Response to treatment for low severity patients Assessed with: HAM-D Follow up: mean 16 weeks	60 per 100	47 per 100 (29 to 75)	RR, 0.78 (0.48 to 1.25)	63 (1 trial)	Insufficient ^{b, c}	Comparisons limited to paroxetine and CT.
Remission in high severity patients Assessed with: HAM-D Follow up: mean 16 weeks	36 per 100	23 per 100 (11 to 46)	RR, 0.63 (0.31 to 1.29)	82 (1 trial)	Insufficient ^{b, c}	Comparisons limited to paroxetine and CT.
Remission in low severity patients Assessed with: HAM-D Follow up: mean 16 weeks	50 per 100	33 per 100 (18 to 60)	RR, 0.65 (0.35 to 1.2)	63 (1 trial)	Insufficient ^{b, c}	Comparisons limited to paroxetine and CT.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% confidence interval).

^b Downgraded for imprecision: single study, small sample size, unable to estimate an effect.

^c Downgraded for risk of bias: high attrition and small sample size

CI = confidence interval; CT, cognitive therapy; HAM-D = Hamilton Depression Rating Scale; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 17. Benefits of second-generation antidepressants compared with integrative therapies as a function of severity

Outcomes	Anticipated absolute effects^a (95% CI): <i>Benefit with integrative therapies monotherapy</i>	Anticipated absolute effects^a (95% CI): <i>Benefit with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Response to treatment for high severity patients	NA	NA	NA	0 (0 trials)	Insufficient	None
Response to treatment for low severity patients	NA	NA	NA	0 (0 trials)	Insufficient	None
Remission in high severity patients Assessed with: HAM-D Follow up: mean 8 weeks	45 per 100	40 per 100 (26 to 62)	RR, 0.89 (0.58 to 1.37)	111 (1 trial)	Insufficient ^{b,c}	Comparisons limited to sertraline or citalopram and IPT.
Remission in low severity patients Assessed with: HAM-D Follow up: mean 8 weeks	75 per 100	56 per 100 (43 to 71)	RR, 0.75 (0.58 to 0.96)	153 (1 trial)	Insufficient ^{b,c}	Comparisons limited to sertraline or citalopram and IPT.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% confidence interval).

^b Downgraded for imprecision: single study, small sample size, unable to estimate an effect

^c Downgraded for reporting bias: Selective non-reporting of several outcomes measured at 2 months, including Geriatric Depression Scale scores, QOL, and satisfaction.

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; IPT = interpersonal therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 18. Benefits of second-generation antidepressants compared with third-wave cognitive behavioral therapy monotherapy as a function of severity

Outcomes	Anticipated absolute effects ^a (95% CI): <i>Benefit with third wave cognitive behavioral therapy monotherapy</i>	Anticipated absolute effects ^a (95% CI): <i>Benefit with SGA</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Response to treatment for high severity patients Assessed with: HAM-D Follow up: mean 16 weeks	60 per 100	40 per 100 (26 to 63)	RR, 0.67 (0.43 to 1.05)	82 (1 trial)	Insufficient ^{b, c}	Comparison limited to paroxetine and BA.
Response to treatment for low severity patients Assessed with: HAM-D Follow up: mean 16 weeks	39 per 100	47 per 100 (24 to 90)	RR, 1.5 (0.62 to 2.32)	61 (1 trial)	Insufficient ^{b, c}	Comparison limited to paroxetine and BA.
Remission in high severity patients Assessed with: HAM-D Follow up: mean 16 weeks	36 per 100	17 per 100 (9 to 31)	RR, 0.47 (0.26 to 0.87)	82 (1 trial)	Insufficient ^{b, c}	Comparison limited to paroxetine and BA.
Remission in low severity patients Assessed with: HAM-D Follow up: mean 16 weeks	44 per 100	37 per 100 (18 to 76)	RR, 0.84 (0.41 to 1.72)	61 (1 trial)	Insufficient ^{b, c}	Comparison limited to paroxetine and BA.
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	NR
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	NR

^a The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% confidence interval).

^b Downgraded for imprecision; single study, small sample size, unable to estimate an effect

^c Downgraded for risk of bias: high attrition

CI = confidence interval; BA = behavioral activation therapy; HAM-D = Hamilton Depression Rating Scale; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 19. Benefits of second-generation antidepressants compared with SAmE as a function of baseline depressive severity

Outcomes	Anticipated Absolute Effects ^a): <i>Benefit with SAmE</i>	Anticipated Absolute Effects ^a (95% CI): <i>Benefit with SGA</i>	Impact of severity as an effect modifier	Number of Participants (Trials)	Strength of Evidence	Comments
Response – change in HAM-D score Assessed with: HAM-D Follow up mean 12 weeks	The mean change in HAM-D score in the control group was 6.19	The mean change in HAM-D score in the intervention group was 6.31 Absolute mean difference was 0.21 higher.	No statistically significant interaction between baseline HAM-D score and treatment groups for reduction in HAM-D scores over time (p=0.87).	129 (1 trial)	Insufficient ^{b, c}	Direct evidence is limited to a single study of SAmE versus escitalopram.
Remission	NA	NA	NA	0 (0 trials)	Insufficient	None
Quality of life	NA	NA	NA	0 (0 trials)	Insufficient	None
Functional capacity	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The risk in the intervention group (and its 95% CI) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for high risk of bias: high dropout

^c Downgraded for imprecision: small sample size in a single study

CI = confidence interval; HAM-D = Hamilton Depression Rating Scale; NR: not reported; RR: Risk ratio; SAmE = S-Adenosyl methionine; SGA = second-generation antidepressant

KQ 2: Summary of Findings Tables

Table 20. Benefits of SGA switches compared with other SGA switches for MDD in adults not responding to an initial adequate SGA treatment attempt

Outcomes	Anticipated absolute effects ^a : <i>Benefits with other SGA switches</i>	Anticipated absolute effects ^a (95% CI): <i>Benefits with SGA switches</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Response Assessed with HAM-D-17 or QIDS-SR-16 Followup: 12 to 14 weeks	NA	NA	RR, 0.96 (0.71 to 1.30) ^{e, f}	727 (1 trial)	Moderate ^b	Comparisons limited to switch strategies to bupropion vs. sertraline.
	NA	NA	RR, 0.91 (0.68 to 1.22) ^{e, f}	727 (1 trial)	Moderate ^b	Comparisons limited to switch strategies to bupropion vs. venlafaxine.
	NA	NA	RR, 0.95 (0.71 to 1.26) ^{e, f}	727 (1 trial)	Moderate ^b	Comparisons limited to switch strategies: sertraline vs. venlafaxine.
Remission Assessed with HAM-D-17 or QIDS-SR-16 Followup: 14 weeks	NR	NR	RR, 1.21 (0.84 to 1.75) ^{e, f}	727 (1 trial)	Low ^{c, d}	Comparisons limited to bupropion vs. sertraline vs. venlafaxine switch strategies. No statistically significant differences between any of the individual switch strategies regardless of the measure used.
Mean change in HAM-D score from baseline Followup: 14 weeks	63 per 100	57 per 100 (49 to 68)	RR, 0.91 (0.78 to 1.07) ^f	406 (1 trial)	Low ^{c, d}	Comparison limited to venlafaxine vs. citalopram switch strategies

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for risk of bias: less than 80% of sample provided outcomes at study completion; medication options not all maximized.

^c Downgraded for risk of bias: less than 80% of sample provided outcomes at study completion (~50% did); medication options not all maximized.

^d Downgraded for imprecision: few events reported.

^e Relative Risk as reported in the article.

^f Crude RR

CGI-S = ;CI = confidence interval; CT = cognitive therapy; HAM-D-17 = Hamilton Depression Scale – 17; MADRS = :NR = not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 21. Benefits of SGA switches compared with nonpharmacologic switches for MDD in adults not responding to an initial adequate SGA treatment attempt

Outcomes	Anticipated absolute effects ^a : <i>Benefits with nonpharmacologic switches</i>	Anticipated absolute effects ^a (95% CI): <i>Benefits with SGA switches</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Response Assessed with QIDS-SR-16 Followup: 12 to 14 weeks	22 per 100	27 to 100 (13 to 54)	RR, 1.2 (0.6 to 2.43) ^e	122 (1 trial) ^b	Low ^{c,d}	Comparison limited to SGA (sertraline, bupropion, or venlafaxine) vs. CT switch strategies.
Remission Assessed with HAM-D-17 or QIDS-SR-16 Followup: 14 weeks	25 per 100	28 to 100 (15 to 54)	RR, 1.12 (0.58 to 2.16) ^{b,e}	122 (1 trial) ^b	Low ^{c,d}	Comparison limited to SGA (sertraline, bupropion, or venlafaxine) vs. CT switch strategies.
Mean change in HAM-D score from baseline Followup: 14 weeks	NR	NR	Not estimable	122 (1 trial) ^b	Low ^{c,d}	Comparison limited to SGA (sertraline, bupropion, or venlafaxine) vs. CT switch strategies.

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b QIDS-SR-16 remission rates led to similar conclusions as those measured by HAM-D-17: RR (95% CI) = 0.88 (0.48 to 1.60).

^c Downgraded for risk of bias: less than 80% of sample provided outcomes at study completion (~50% did); medication options not all maximized.

^d Downgraded for imprecision: single study.

^e Crude RR

CI = confidence interval; CT = cognitive therapy; HAM-D-17 = Hamilton Depression Scale – 17; NR = not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 22. Benefits of SGA augmentation compared with SGA augmentation for MDD in adults not responding to an initial adequate SGA treatment attempt

Outcomes	Anticipated absolute effects ^a : <i>Benefits with SGA augmentation</i>	Anticipated absolute effects ^a (95% CI): <i>Benefits with SGA augmentation</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Response Assessed with QIDS-SR-16 Followup:14 weeks	27 per 100	32 to 100 (25 to 41)	RR, 1.18 (0.92 to 1.53) ^e	565 (1 trial)	Low ^{c, d}	Comparison limited to bupropion vs. buspirone augmentation of citalopram treatment.
Remission Assessed with HAM-D-17 or QIDS-SR-16 Followup:14 weeks	30 per 100	30 to 100 (23 to 38)	RR, 0.99 (0.77 to 1.27) ^{b, e}	565 (1 trial)	Low ^{c, d}	Comparison limited to bupropion vs. buspirone augmentation of citalopram treatment
Mean change in QIDS-SR-16 score from baseline Followup:14 weeks	NR	NR	Not estimable	565 (1 trial)	Low ^{c, d}	Comparison limited to bupropion vs. buspirone augmentation of citalopram treatment.

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b QIDS-SR-16 data did not affect conclusions about comparative remission rates: RR (95% CI) = 1.19 (0.95 to 1.48).

^c Downgraded for risk of bias: less than 80% of sample provided outcomes at study completion (~50% did); medication options not all maximized.

^d Downgraded for precision: few events reported.

^e Crude RR

CI = confidence interval; CT = cognitive therapy; HAM-D-17 = Hamilton Depression Scale – 17; NR = not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 23. Benefits of SGA augmentation compared with nonpharmacologic augmentation for MDD in adults not responding to an initial adequate SGA treatment attempt

Outcomes	Anticipated absolute effects ^a : <i>Benefits with nonpharmacologic augmentation</i>	Anticipated absolute effects ^a (95% CI): <i>Benefits with SGA augmentation</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Response Assessed with QIDS-SR-16 Followup: 14 weeks	35 per 100	28 to 100 (18 to 43)	RR, 0.8 (0.51 to 1.23) ^e	182 (1 trial)	Low ^{b, c}	Comparison limited to SGA (bupropion or buspirone) vs. CT augmentation of citalopram treatment.
Remission Assessed with HAM-D-17 or QIDS-SR-16 Followup: 14 weeks	23 per 100	33 to 100 (20 to 56)	RR, 1.44 (0.87 to 2.41) ^{b, e}	182 (1 trial)	Low ^{b, c}	Comparison limited to SGA (bupropion or buspirone) vs. CT augmentation of citalopram treatment.
Mean change in QIDS-SR-16 score from baseline Followup: 14 weeks	NR	NR	Not estimable	182 (1 trial)	Low ^{b, c}	Comparison limited to SGA (bupropion or buspirone) vs. CT augmentation of citalopram treatment.

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b QIDS-SR-16 data did not affect conclusions about comparative remission rates: RR (95% CI) = 1.08 (0.69 to 1.69).

^c Downgraded for risk of bias: less than 80% of sample provided outcomes at study completion (~50% did); medication options not all maximized.

^d Downgraded for imprecision: few events reported.

^e Crude RR

CI = confidence interval; CT = cognitive therapy; HAM-D-17 = Hamilton Depression Scale – 17; NR = not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 24. Benefit of SGA switches for MDD in adults not responding to an initial adequate SGA treatment attempt as a function of baseline severity

Outcomes	Anticipated absolute effects^a: <i>Benefit with higher severity</i>	Anticipated absolute effects^a (95% CI): <i>Benefit with lower severity</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Response	NR	NR	Not estimable	0 (0 trials)	Insufficient	None
Remission Assessed with HAM-D-17 Followup: 12 to 14 weeks	NR	NR	Not estimable	(1 trials) ^b	Insufficient ^{c, d}	Comparisons limited to venlafaxine vs. citalopram switch strategies or to bupropion vs. sertraline vs. venlafaxine switch strategies.
Mean change in QIDS-SR-16 score from baseline	NR	NR	Not estimable	0 (0 trials)	Insufficient	None

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Two secondary analyses of two different RCTs

^c Downgraded for risk of bias: selective outcome reporting bias in one study

^d Downgraded for inconsistency: two studies reported contrasting results

CI = confidence interval; CT = cognitive therapy; HAM-D-17 = Hamilton Depression Scale – 17; NR = not reported; QIDS-SR-16 = ;RR = risk ratio; SGA = second-generation antidepressant

KQ 3 and KQ4: Summary of Findings Tables

Table 25. Risks of SGAs compared with CBT for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with CBT</i>	Anticipated absolute effects ^a (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality Follow up: range 8 to 16 weeks	1 per 100	2 per 100 (0 to 8)	RR, 1.2 (0.24 to 6.13)	465 (4 trials) ^k	Insufficient ^{ff, h}	None
Serious adverse events Follow up: range 8 to 10 weeks	NR	NR	RR, 5.44 (0.31 to 96.36)	140 (2 trials) ^k	Insufficient ^{fg, h}	None
Overall risk for overall adverse events: Follow up: mean 13 weeks	1 per 100	16 per 100 (2 to 122)	RR, 17.84 (2.32 to 137.4)	170 (1 trial)	Insufficient ^{bd}	Results for SGAs appear to substantially underestimate the risk of adverse events. A comprehensive systematic assessments of the risk of harms for SGAs reported that, on average, 60 percent of patients treated with SGAs experience at least one adverse event during the course of treatment.
Overall discontinuation Follow up: range 8 to 14 weeks	17 per 100	14 per 100 (10 to 21)	RR, 0.85 (0.55 to 1.30)	560 (4 trials) ⁱ	Moderate ^f	Second-generation antidepressants are limited to fluoxetine, fluvoxamine, and paroxetine.
Discontinuation because of adverse events Follow up: range 8 to 14 weeks	2 per 100	8 per 100 (3 to 23)	RR, 2.00 (0.74 to 5.45)	388 (3 trials) ^j	Low ^{ce}	Discontinuation rates because of adverse events were statistically significantly higher for patients on SGAs than on CBT when we included two high risk of bias studies (RR 3.09, 96% CI 1.14-8.35)

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: very few events; 95% confidence intervals wide.

^c Not upgraded for large effect because of extreme imprecision.

^d Downgraded for risk of bias: adverse events reported only for study completers.

^e Downgraded for imprecision: very few events; 95% confidence intervals especially wide and cross both thresholds of appreciable differences

^f Downgraded for imprecision: very few events

^g Downgraded for risk of bias: one study's data for subset of patients with MDD not based on ITT analysis

^h Downgraded for risk of bias: two of 4 studies had a high risk of bias due to high overall and differential attrition rates; in one study, attrition particularly high among SGA patients; in another study, available data based on completers analysis only

ⁱ Does not include data from 4 high risk of bias studies because sensitivity analysis including those studies did not change meta-analysis findings

^j Does not include data from 2 high risk of bias studies because sensitivity analysis including those studies did not change meta-analysis findings

^k Includes high risk of bias evidence because number of studies with lower risk of bias insufficient to allow meta-analysis of findings

CBT = cognitive behavioral therapy; CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 26. Risks of SGAs compared with combinations of SGA and CBT for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with combination SGA and CBT</i>	Anticipated absolute effects ^a : (95% CI): <i>Risk with SGA switches</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality	NA	NA	NA	0 (0 trials)	Insufficient	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Overall discontinuation Follow up: mean 16 weeks	16 per 100	12 per 100 (6 to 26)	RR, 0.77 (0.37 to 1.6)	176 (2 trials)	Low ^b	Comparison limited to escitalopram with escitalopram combined with telephone CBT
Discontinuation because of adverse events Follow up: mean 12 weeks	2 per 100	8 per 100 (2 to 37)	RR, 3.42 (0.73 to 16.01)	176 (2 trials)	Low ^{b, c}	Comparison limited to escitalopram with escitalopram combined with telephone CBT

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: very few events; very wide 95% confidence interval across both thresholds of appreciable differences.

^c Not upgraded for large effect size because of extreme imprecision.

CBT = cognitive behavioral therapy; CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 27. Risks of SGAs compared with integrative therapies for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with integrative therapies</i>	Anticipated absolute effects ^a (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality Follow up: mean 12 weeks	16 per 100	6 per 100 (3 to 13)	RR, 0.39 (0.19 to 0.82)	291 (2 trials)	Insufficient ^{b, f}	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events	NA	NA	NA	0 (0 trials)	Insufficient	Comparison limited to escitalopram versus interpersonal therapy
Overall discontinuation Follow up: mean 13 weeks	11 per 100	16 per 100 (10 to 26)	RR, 1.38 (0.83 to 2.3)	384 (2 trials)	Insufficient ^{b, e}	Comparison limited to citalopram, nefazodone, and sertraline vs. Interpersonal therapy
Discontinuation because of adverse events Follow up: mean 13 weeks	NR	NR	RR, 0.33 (0.01 to 8.06)	287 (1 trial)	Insufficient ^{b, d}	Comparison limited to citalopram and sertraline versus interpersonal therapy

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: very few events.

^c Downgraded for risk of bias: high attrition rate; unclear whether outcome assessors were masked; in one study, no indication that suicidality incidence data adjusted for baseline presence of suicidality or that ITT analysis applied to suicidality data.

^d Downgraded for risk of bias: outcomes reporting bias; most studies did not report on discontinuation because of adverse events

^e Downgraded for risk of bias: one of two available studies did not report discontinuations taking place between randomization and onset of treatment; impossible to determine how unreported discontinuations would have affected our findings.

^f Downgraded for imprecision: very few events in both studies, and in one study, 95% confidence interval crosses both thresholds of appreciable differences.

CI = confidence interval; CT = cognitive therapy; NR = not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 28. Risks of SGAs compared with combination of SGAs and integrative therapies for MDD in adults

Outcomes	Anticipated absolute effects^a: <i>Risk with combination of SGAs and integrative therapies</i>	Anticipated absolute effects^a: (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality	NA	NA	NA	0 (0 trials)	Insufficient	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events Follow up: mean 12 weeks	NA	NA	NA	0 (0 trials)	Insufficient	None
Overall discontinuation Follow up: mean 16 weeks	25 per 100	27 per 100 (16 to 47)	RR, 1.11 (0.64 to 1.93)	96 (1 trial)	Insufficient ^{b, c}	Comparison limited to one trial of nefazodone with a combination of nefazodone and integrative therapy
Discontinuation because of adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None

^aThe benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: very few events; very wide 95% confidence interval across both thresholds of appreciable differences.

^c Not upgraded for large effect size because of extreme imprecision.

CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 29. Risks of SGAs compared with psychodynamic therapy for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with psychodynamic therapy</i>	Anticipated absolute effects ^a : (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality Follow up: mean 8 weeks	16 per 100	16 per 100 (7 to 34)	RR, 1.01 (0.47 to 2.09)	141 (1 trial)	Insufficient ^{b, e}	None
Suicidality Follow up: mean 96 weeks	3 per 100	4 per 100 (1 to 19)	RR, 1.32 (0.3 to 5.73)	181 (1 trial)	Low ^b	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Overall discontinuation Follow up: mean 16 weeks	25 per 100	17 per 100 (10 to 30)	RR, 0.68 (0.39 to 1.2)	192 (2 trials)	Insufficient ^{b, c, d}	Comparisons are limited to fluoxetine and venlafaxine vs. psychodynamic therapy
Overall discontinuation Follow up: mean 48 weeks	19 per 100	24 per 100 (9 to 69)	RR, 1.25 (0.44 to 3.57)	51 (1 trial)	Low ^b	None
Overall discontinuation Follow up: mean 96 weeks	19 per 100	15 per 100 (8 to 29)	RR, 0.81 (0.43 to 1.55)	181 (1 trial)	Low ^b	None
Discontinuation because of adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: very few events; 95% confidence intervals cross both thresholds of appreciable differences.

^c Downgraded for risk of bias: outcomes reporting bias in one of two available studies, in that attrition taking place between randomization and treatment onset left unreported.

^d Downgraded for inconsistency: conflicting directionality of point estimates in the two available studies for overall discontinuation.

^e Downgraded for risk of bias: high overall attrition and unclear how that attrition affected incidence rates of suicidality, despite use of modified ITT analysis.

CI = confidence interval; CT = cognitive therapy; NR = not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 30. Risks of SGAs compared with combination of SGAs and psychodynamic therapy for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with combination of SGAs and psychodynamic therapy</i>	Anticipated absolute effects ^a : (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality Follow up: mean 96 weeks	1 per 100	4 per 100 (1 to 39)	RR, 4.00 (0.46 to 35.1)	182 (1 trial)	Low ^{b, c}	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Overall discontinuation Follow up: mean 96 weeks	32 per 100	15 per 100 (9 to 27)	RR, 0.48 (0.27 to 0.85)	182 (1 trial)	Low ^b	Comparison is limited to fluoxetine with a combination of fluoxetine and long-term psychodynamic therapy
Discontinuation because of adverse events	0 per 100	0 to 100 (0 to 0)	Not estimable	0 (0 trials)	Insufficient	None

^aThe benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecisions: very few events; confidence intervals crosses both thresholds of appreciable differences.

^c Not upgraded for large effect because of extreme imprecision.

CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 31. Risks of SGA compared with third wave CBT for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with third wave CBT</i>	Anticipated absolute effects ^a (95% CI): <i>Risk with SGA switches</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality Follow up: mean 16 weeks	NA	NA	NA	143(1 trial)	Insufficient ^{b, c, f}	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events Follow up: mean 13 weeks	NA	NA	NA	0 (0 trials)	Insufficient	None
Overall discontinuation Follow up: mean 13 weeks	10 per 100	27 per 100 (14 to 52)	RR, 2.76 (1.4 to 5.41)	243 (2 trials)	Insufficient ^{b, c, d}	None
Discontinuation because of adverse events Follow up: mean 13 weeks	3 per 100	23 per 100 (1 to 43)	RR, 7.00 (0.37 to 132.11)	243 (2 trials)	Insufficient ^{b, c, e}	None

^aThe benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: very few events; 95% confidence intervals wide

^c Not upgraded for large effect because of extreme imprecision

^d Downgraded for risk of bias: in one of 2 available studies, high differential attrition rate between SGA and third-wave CBT groups may have been affected by statistically significant difference in percentages of women assigned to each treatment

^e Downgraded for imprecision: very few events; 95% confidence intervals very wide and cross both thresholds of appreciable differences

^f Downgraded for risk of bias: in only available study reporting suicidality data, statistically significant baseline difference in percentages of women assigned to each treatment may have artificially affected risk of suicidality

CBT = cognitive behavioral therapy; CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 32. Risks of SGAs compared with any psychological therapy for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with any psychological therapy</i>	Anticipated absolute effects ^a (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality Follow up: range 8 to 16 weeks	9 per 100	8 per 100 (4 to 17)	RR, 0.87 (0.42 to 1.77)	1227 (8 trials)	Low ^{fb, c}	None
Serious adverse events Follow up: mean 8 weeks	NR	NR	RR, 4.87 (0.27 to 87.69)	92 (1 trial)	Low ^{fb, ?}	None
Overall risk for overall adverse events: Follow up: mean 14 weeks	1 per 100	16 per 100 (2 to 122)	RR, 17.84 (2.32 to 137.4)	170 (1 trial)	Insufficient ^{b, c}	None
Overall discontinuation Follow up: range 8 to 16 weeks	16 per 100	19 per 100 (15 to 23)	RR, 1.23 (0.79 to 1.91)	1679 (7 trials)	Moderate ^b	Interventions are limited to: 1) fluoxetine, fluvoxamine, paroxetine, sertraline, and 2) behavioral activation, cognitive therapy, problem solving therapy, rational emotive behavior therapy, short-term psychodynamic supportive psychotherapy
Discontinuation because of adverse events Follow up: range 8 to 16 weeks	2 per 100	7 per 100 (4 to 15)	RR, 2.64 (0.98 to 7.08)	1097 (7 trials)	Moderate ^b	Interventions are limited to: 1) fluoxetine, fluvoxamine, paroxetine, sertraline, and 2) behavioral activation, cognitive therapy, problem solving therapy, rational emotive

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: few events.

^c Downgraded for risk of bias: very high attrition in two studies.

^d Downgraded for risk of bias: adverse events reported only for study completers.

CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 33. Risks of SGAs compared with acupuncture for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with acupuncture</i>	Anticipated absolute effects ^a : (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality	NA	NA	NA	0 (0 trials)	Insufficient	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Overall risk for overall adverse events: direct evidence Follow up: mean 6 weeks	6 per 100	4 per 100 (1 to 24)	RR, 0.69 (0.12 to 3.98)	98 (1 trial)	Insufficient ^{f,g}	None
Overall risk for overall adverse events: indirect evidence Follow up: mean 8 weeks	10 per 100	40 per 100 (35 to 47)	RR, 3.96 (3.4 to 4.62)	3128 (21 trials)	Moderate ^b	A systematic review which did not meet our eligibility criteria because it also included other depressive disorders than MDD provides the most comprehensive assessment of the comparative risk of harms between SGAs and acupuncture
Overall discontinuation Follow up: mean 6 weeks	56 per 100	2 per 100 (0 to 31)	RR, 0.03 (0 to 0.56)	75 (1 trial)	Low ^{c,d,e}	None
Discontinuation because of adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for indirectness: numbers are based on a systematic review that included all depressive disorders and some first generation antidepressants.

^c Downgraded for imprecision: very few events; small sample size well below optimal information size.

^d Not downgraded for risk of bias because lack of ITT analysis does not affect the validity of discontinuation outcomes.

^e Not upgraded for large effect because of extreme imprecision.

^f Downgraded for risk of bias: validity of data in question due to lack of reporting about key components of study design, including randomization, allocation concealment, between-group similarity of baseline characteristics, and use of blinded outcome assessment.

^g Downgraded for imprecision: very few events; 95% confidence interval crosses both thresholds of appreciable differences.

CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 34. Risks of SGAs compared with combination of SGAs and acupuncture for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with combination of SGA and acupuncture</i>	Anticipated absolute effects ^a : (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality Follow up: mean 8 weeks	NR	NR	Not estimable	0 (0 trials)	Insufficient	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events Follow up: mean 8 weeks	4 per 100	4 per 100 (1 to 21)	RR, 1.00 (0.21 to 4.79)	140 (1 trial)	Low ^b	None
Overall discontinuation Follow up: mean 6 weeks	10 per 100	13 per 100 (5 to 31)	RR, 1.29 (0.53 to 3.14)	428 (3 trials)	Moderate ^c	None
Discontinuation because of adverse events Follow up: mean 6 weeks	2 per 100	1 per 100 (0 to 7)	RR, 0.37 (0.04 to 3.54)	288 (2 trials)	Low ^b	None

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: very few events; 95% confidence interval crosses both thresholds of appreciable differences.

^c Downgraded for imprecision: few events; 95% confidence interval nearly crosses both thresholds of appreciable differences; not downgraded any further because overall sample size exceeds optimal information size of 300.

^d Crude RR

CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 35. Risks of SGAs compared with omega-3 fatty acids for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with omega-3 fatty acids</i>	Anticipated absolute effects ^a : (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality Follow up: mean 8 weeks	5 per 100	2 per 100 (0 to 39)	RR, 0.33 (0.01 to 7.72)	40 (1 trial)	Insufficient ^{b, c}	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Overall discontinuation Follow up: mean 4 weeks	15 per 100	15 per 100 (4 to 66)	RR, 1.0 (0.23 to 4.37)	40 (1 trial)	Low ^b	None
Discontinuation because of adverse events Follow up: mean 8 weeks	5 per 100	5 per 100 (0 to 75)	RR, 1.0 (0.07 to 14.9)	40 (1 trial)	Low ^b	None

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: very few events; 95% confidence intervals cross both thresholds of appreciable differences.

^c Downgraded for risk of bias: high drop out rates, no intention-to-treat analyses available.

CI = confidence interval; CT = cognitive therapy; NR = not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 36. Risks of SGAs compared with combination of SGAs and omega-3 fatty acids for MDD in adults

Outcomes	Anticipated absolute effects^a: <i>Risk with combination of SGAs and omega-3 fatty acids</i>	Anticipated absolute effects^a: (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality	NA	NA	NA	0 (0 trials)	Insufficient	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events Follow up: mean 12 weeks	NA	NA	NA	0 (0 trials)	Insufficient	None
Overall discontinuation Follow up: mean 4 weeks	10 per 100	24 per 100 (8 to 70)	RR, 2.38 (0.81 to 6.98)	82 (2 trials)	Low ^{c, d}	Overall discontinuation rates were also similar between fluoxetine and a combination of fluoxetine and omega-3 fatty acids
Discontinuation because of adverse events Follow up: mean 8 weeks	10 per 100	5 per 100 (.5 to 51)	RR, 0.05 (0.05 to 508)	40 (1 trial)	Low ^b	None

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: very few events; 95% confidence interval crosses both thresholds of appreciable differences.

^c Not upgraded for large effect because of imprecision

^d Downgraded for imprecision: very few events; 95% confidence intervals nearly cross both thresholds of appreciable differences.

CI = confidence interval; CT = cognitive therapy; NR = not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 37. Risks of SGAs compared with SAmE for MDD in adults

Outcomes	Anticipated absolute effects^a: <i>Risk with SAmE</i>	Anticipated absolute effects^a (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality Follow up: mean 8 weeks	NA	NA	NA	0 (0 trials)	Insufficient	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events	NA	NA	NA	0 (0 trials)	Insufficient	No information on overall risk of adverse events. One study provided data about selected adverse events only.
Overall discontinuation Follow up: mean 12 weeks	44 per 100	52 per 100 (34 to 79)	RR, 1.19 (0.78 to 1.8)	129 (1 trial)	Low ^d	None
Discontinuation because of adverse events Follow up: mean 12 weeks	5 per 100	12 per 100 (3 to 44)	RR, 2.63 (0.73 to 9.46)	129 (1 trial)	Insufficient ^{b, c}	None

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: few events; 95% confidence interval crosses both thresholds of appreciable differences.

^c Downgraded for risk of bias: very high overall drop out rate.

^d Downgraded for imprecision: few events; 95% confidence interval crosses nearly both thresholds of appreciable differences

CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 38. Risks of SGAs compared with St. John's wort for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with St. John's wort</i>	Anticipated absolute effects ^a : (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality	1 per 100	1 per 100 (0 to 10)	RR, 1.03 (0.07 to 16.34)	331 (3 trials)	Insufficient ^{h, k}	None
Serious adverse events	2 per 100	1 per 100 (0 to 3)	RR, 0.63 (0.21 to 1.93)	932 (5 trials)	Low ^{f, g}	None
Risk for overall adverse events	39 per 100	47 per 100 (41 to 53)	RR, 1.19 (1.05 to 1.34)	1427 (8 trials)	Moderate ^{d, e}	None
Risk for overall adverse events in subgroup based on older age	65 per 100	84 [rt 100	RR, 1.30 (0.66 to 2.54)	131 (1 trial)	Low ⁱ	Comparison limited to fluoxetine and SJW in older adults (60 to 80 years)
Overall discontinuation	14 per 100	18 per 100 (14 to 22)	RR, 1.29 (1.04 to 1.6)	1813 (12 trials)	Moderate ^{b, c}	None
Discontinuation because of adverse events	4 per 100	7 per 100 (5 to 10)	RR, 1.79 (1.19 to 2.71)	1773 (11 trials)	Moderate ^{h, i}	Crude RR
Discontinuation because of adverse events in subgroup based on older age	65 per 100	79 per 100 (29 to 218)	RR, 1.22 (0.44 to 3.36)	161 (1 trial)	Low ^j	Comparison limited to fluoxetine and SJW in older adults (60 to 80 years)

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Three of 12 included trials were rated as having a high risk of bias because of how they reported harms data, including overall discontinuation. In addition, one study with a low risk of bias rating reported conflicting discontinuation data, making it unclear which of its data were accurate. Still, sensitivity analysis results no different than primary analysis.

^c <300 events overall (N=281), and summary point estimate non-significant, but 95% CI crosses the 1.25 threshold.

^d Substantial heterogeneity (I-squared = 65%) in the directionality of point estimates, although all but two trials' 95% CIs overlapped. Specifically, three of 8 included trials found that patients taking SJW experienced fewer overall adverse events, in contrast to what the other five trials found.

^e Summary point estimate non-significant, and its 95% CI crosses the 1.25 threshold.

^f Summary point estimate non-significant, but its 95% CI crosses both the 0.75 and 1.25 thresholds. Also, far fewer than 300 events overall (N=13).

^g One of five included trials was rated as having a high risk of bias.

^h Two of 11 included trials were rated as having a high risk of bias.

ⁱ <300 events overall (N=94).

^j Only 2 events overall, 1 per relevant study.

^k Conflicting evidence and no clear pattern emerging because only one case of suicidality in each treatment group.

^l Downgraded for serious imprecision; does not fulfill OIS, small number of events

CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 39. Risks of SGAs compared with exercise for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with exercise</i>	Anticipated absolute effects ^a (95% CI): <i>Risk with SGAs</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality	NA	NA	NA	0 (0 trials)	Insufficient	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Overall discontinuation Follow up: mean 16 weeks	2 per 100	1 per 100 (1 to 3)	RR, 0.87 (0.48 to 1.59)	254 (2 trials)	Low ^b	None
Discontinuation because of adverse events Follow up: mean 16 weeks	NR	NR	RR, 20.96 (1.19 to 367.97)	254 (2 trials)	Low ^{c, d}	Comparison limited to sertraline vs. exercise. Patients treated with a combination of sertraline and exercise had similar discontinuation rates because of adverse events as patients on sertraline monotherapy (9% vs. 10%).

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: few events; confidence interval crosses threshold of appreciable difference.

^c Downgraded for imprecision: very few events; very wide confidence interval.

^d Not upgraded for large effect size because of extreme imprecision.

CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 40. Risks of SGAs compared with combination of SGA and exercise for MDD in adults

Outcomes	Anticipated absolute effects ^a : Risk with SGA and exercise	Anticipated absolute effects ^a (95% CI): Risk with SGAs	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality	NA	NA	NA	0 (0 trials)	Insufficient	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events Follow up: mean 12 weeks	NA	NA	NA	0 (0 trials)	Insufficient	None
Overall discontinuation Follow up: mean 16 weeks	20 per 100	15 per 100 (6 to 35)	RR, 0.73 (0.31 to 1.73)	103 (1 trial)	Low ^b	None
Discontinuation because of adverse events Follow up: mean 16 weeks	9 per 100	10 per 100 (3 to 34)	RR, 1.15 (0.35 to 3.72)	103 (1 trial)	Low ^b	Comparison limited to sertraline vs. Sertraline + exercise

^aThe benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: very few events; very wide 95% confidence interval across both thresholds of appreciable differences.

CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Table 41. Risks of SGA switches compared with other SGA switches for MDD in adults

Outcomes	Anticipated absolute effects ^a : <i>Risk with other SGA switches</i>	Anticipated absolute effects ^a : (95% CI): <i>Risk with SGA switches</i>	Relative effect (95% CI)	Number of participants (Trials)	Strength of evidence	Comments
Suicidality	NA	NA	NA	0 (0 trials)	Insufficient	None
Serious adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None
Risk for overall adverse events Follow up: mean 12 weeks	63 per 100	57 per 100 (49 to 68)	RR, 0.91 (0.78 to 1.07)	406 (1 trial)	Low ^{b,c}	Comparison limited to switch to venlafaxine vs. switch to citalopram
Overall discontinuation Follow up: mean 12 weeks	21 per 100	24 per 100 (17 to 35)	RR, 1.17 (0.82 to 1.68)	406 (1 trial)	Low ^{b,c}	Comparison limited to switch to venlafaxine vs. switch to citalopram
Discontinuation because of adverse events	NA	NA	NA	0 (0 trials)	Insufficient	None

^a The benefit or risk in the intervention group (and its 95% confidence interval) is based on the assumed benefit or risk in the comparison group and the relative effect of the intervention (and its 95% CI).

^b Downgraded for imprecision: small number of events.

^c Downgraded for risk of bias: potential confounding from prior treatment attempts with psychotherapy, which was not accounted for at baseline.

CI = confidence interval; CT = cognitive therapy; NR: not reported; RR = risk ratio; SGA = second-generation antidepressant

Appendix F. Studies Included in Network Meta-Analyses

1. Institute for Quality and Efficiency in Health Care (IQWiG). Selektive Serotonin und Noradrenalin Wiederaufnahmehemmer (SNRI) bei Patienten mit Depressionen, A05-20A, Version 1.0. Institute for Quality and Efficiency in Health Care (IQWiG) (c) IQWiG (Institute for Quality and Efficiency in Health Care). Cologne, Germany: 2009.
2. Allen JJ, Schnyer RN, Chambers AS, et al. Acupuncture for depression: a randomized controlled trial. *J Clin Psychiatry*. 2006 Nov;67(11):1665-73. Epub: 2007/01/02. PMID: 17196044.
3. Allen JJB, Schnyer RN, Hitt SK. The efficacy of acupuncture in the treatment of major depression in women. *Psychol Sci*. 1998;9(5):397-401. PMID: 1998-11225-010. First Author & Affiliation: Allen, John J. B.
4. Alvarez E, Perez V, Dragheim M, et al. A double-blind, randomized, placebo-controlled, active reference study of Lu AA21004 in patients with major depressive disorder. *Int J Neuropsychopharmacol*. 2012 Jun;15(5):589-600. Epub: 2011/07/20. PMID: 21767441.
5. Alves C, Cachola I, Brandao J. Efficacy and tolerability of venlafaxine and fluoxetine in outpatients with major depression. *Primary Care Psychiatry*. 1999;5(2):57-63.
6. Bakish D, Bose A, Gommoll C, et al. Levomilnacipran ER 40 mg and 80 mg in patients with major depressive disorder: a phase III, randomized, double-blind, fixed-dose, placebo-controlled study. *J Psychiatry Neurosci*. 2014 Jan;39(1):40-9. Epub: 2013/10/23. PMID: 24144196.
7. Barber JP, Barrett MS, Gallop R, et al. Short-term dynamic psychotherapy versus pharmacotherapy for major depressive disorder: a randomized, placebo-controlled trial. *J Clin Psychiatry*. 2012 Jan;73(1):66-73. Epub: 2011/12/14. PMID: 22152401.
8. Beasley CM, Dornseif BE, Pultz JA, et al. Fluoxetine versus trazodone: efficacy and activating-sedating effects. *J Clin Psychiatry*. 1991;52(7):294-9. PMID: CN-00076754.
9. Behnke K, Jensen GS, Graubaum HJ, et al. Hypericum perforatum versus fluoxetine in the treatment of mild to moderate depression. *Adv Ther*. 2002 Jan-Feb;19(1):43-52. Epub: 2002/05/15. PMID: 12008860.
10. Benkert O, Szegedi A, Kohlen R. Mirtazapine compared with paroxetine in major depression. *J Clin Psychiatry*. 2000 Sep;61(9):656-63. Epub: 2000/10/13. PMID: 11030486.
11. Bielski RJ, Ventura D, Chang CC. A double-blind comparison of escitalopram and venlafaxine extended release in the treatment of major depressive disorder. *J Clin Psychiatry*. 2004 Sep;65(9):1190-6. Epub: 2004/09/16. PMID: 15367045.
12. Bjerkenstedt L, Edman GV, Alken RG, et al. Hypericum extract LI 160 and fluoxetine in mild to moderate depression: a randomized, placebo-controlled multi-center study in outpatients. *Eur Arch Psychiatry Clin Neurosci*. 2005 Feb;255(1):40-7. Epub: 2004/11/13. PMID: 15538592.
13. Blumenthal JA, Babyak MA, Doraiswamy PM, et al. Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med*. 2007 Sep-Oct;69(7):587-96. Epub: 2007/09/12. PMID: 17846259.
14. Boyer P, Montgomery S, Lepola U, et al. Efficacy, safety, and tolerability of fixed-dose desvenlafaxine 50 and 100 mg/day for major depressive disorder in a placebo-controlled trial. *Int Clin Psychopharmacol*. 2008 Sep;23(5):243-53. Epub: 2008/08/16. PMID: 18703933.
15. Brannan SK, Mallinckrodt CH, Brown EB, et al. Duloxetine 60 mg once-daily in the treatment of painful physical symptoms in patients with major depressive disorder. *J Psychiatr Res*. 2005 Jan;39(1):43-53. Epub: 2004/10/27. PMID: 15504423.
16. Claghorn JL. The safety and efficacy of paroxetine compared with placebo in a double-blind trial of depressed outpatients. *J Clin Psychiatry*. 1992 Feb;53 Suppl:33-5. Epub: 1992/02/01. PMID: 1531821.

17. Cohn CK, Robinson DS, Roberts DL, et al. Responders to antidepressant drug treatment: a study comparing nefazodone, imipramine, and placebo in patients with major depression. *J Clin Psychiatry*. 1996;57 Suppl 2:15-8. Epub: 1996/01/01. PMID: 8626358.
18. Coleman CC, Cunningham LA, Foster VJ, et al. Sexual dysfunction associated with the treatment of depression: a placebo-controlled comparison of bupropion sustained release and sertraline treatment. *Ann Clin Psychiatry*. 1999 Dec;11(4):205-15. Epub: 1999/12/22. PMID: 10596735.
19. Coleman CC, King BR, Bolden-Watson C, et al. A placebo-controlled comparison of the effects on sexual functioning of bupropion sustained release and fluoxetine. *Clin Ther*. 2001 Jul;23(7):1040-58. Epub: 2001/08/25. PMID: 11519769.
20. Company ELA. Duloxetine versus placebo in the treatment of major depression: study F1J-MC-HMAQ, study group B; clinical study summary #3327 2006. p. In: *ClinicalStudyResults*.
21. Company ELA. Duloxetine versus placebo and paroxetine in the acute treatment of major depression: study F1J-MC-HMAT, study group A; clinical study summary #4091. 2006. p. In: *ClinicalStudyResults*.
22. Corrigan MH, Denahan AQ, Wright CE, et al. Comparison of pramipexole, fluoxetine, and placebo in patients with major depression. *Depress Anxiety*. 2000;11(2):58-65. Epub: 2000/05/17. PMID: 10812530.
23. Croft H, Settle E, Jr., Houser T, et al. A placebo-controlled comparison of the antidepressant efficacy and effects on sexual functioning of sustained-release bupropion and sertraline. *Clin Ther*. 1999 Apr;21(4):643-58. Epub: 1999/06/11. PMID: 10363731.
24. De Nayer A, Geerts S, Ruelens L, et al. Venlafaxine compared with fluoxetine in outpatients with depression and concomitant anxiety. *Int J Neuropsychopharmacol*. 2002;5(2):115-20. PMID: 2002-06574-003. PMID: 12135535. First Author & Affiliation: De Nayer, Andre.
25. Detke MJ, Lu Y, Goldstein DJ, et al. Duloxetine, 60 mg once daily, for major depressive disorder: a randomized double-blind placebo-controlled trial. *J Clin Psychiatry*. 2002 Apr;63(4):308-15. Epub: 2002/05/10. PMID: 12000204.
26. Detke MJ, Lu Y, Goldstein DJ, et al. Duloxetine 60 mg once daily dosing versus placebo in the acute treatment of major depression. *J Psychiatr Res*. 2002 Nov-Dec;36(6):383-90. Epub: 2002/10/24. PMID: 12393307.
27. Detke MJ, Wiltse CG, Mallinckrodt CH, et al. Duloxetine in the acute and long-term treatment of major depressive disorder: a placebo- and paroxetine-controlled trial. *Eur Neuropsychopharmacol*. 2004 Dec;14(6):457-70. Epub: 2004/12/14. PMID: 15589385.
28. Dierick M, Ravizza L, Realini R, et al. A double-blind comparison of venlafaxine and fluoxetine for treatment of major depression in outpatients. *Prog Neuropsychopharmacol Biol Psychiatry*. 1996 Jan;20(1):57-71. Epub: 1996/01/01. PMID: 8861177.
29. Dunlop BW, Reddy S, Yang L, et al. Symptomatic and functional improvement in employed depressed patients: a double-blind clinical trial of desvenlafaxine versus placebo. *J Clin Psychopharmacol*. 2011 Oct;31(5):569-76. Epub: 2011/08/27. PMID: 21869698.
30. Dunn AL, Trivedi MH, Kampert JB, et al. Exercise treatment for depression: efficacy and dose response. *Am J Prev Med*. 2005 Jan;28(1):1-8. Epub: 2005/01/01. PMID: 15626549.
31. Fabre L, Birkhimer LJ, Zaborny BA, et al. Fluvoxamine versus imipramine and placebo: a double-blind comparison in depressed patients. *Int Clin Psychopharmacol*. 1996 Jun;11(2):119-27. Epub: 1996/06/01. PMID: 8803649.
32. Fava M, Amsterdam JD, Deltito JA, et al. A double-blind study of paroxetine, fluoxetine, and placebo in outpatients with major depression. *Ann Clin Psychiatry*. 1998 Dec;10(4):145-50. Epub: 1999/02/13. PMID: 9988054.

33. Feiger A, Kiev A, Shrivastava RK, et al. Nefazodone versus sertraline in outpatients with major depression: focus on efficacy, tolerability, and effects on sexual function and satisfaction. *J Clin Psychiatry*. 1996;57 Suppl 2:53-62. Epub: 1996/01/01. PMID: 8626364.
34. Feiger AD, Tourian KA, Rosas GR, et al. A placebo-controlled study evaluating the efficacy and safety of flexible-dose desvenlafaxine treatment in outpatients with major depressive disorder. *CNS Spectr*. 2009 Jan;14(1):41-50. Epub: 2009/01/27. PMID: 19169187.
35. Feighner JP, Boyer WF. Paroxetine in the treatment of depression: a comparison with imipramine and placebo. *J Clin Psychiatry*. 1992 Feb;53 Suppl:44-7. Epub: 1992/02/01. PMID: 1531824.
36. Feighner JP, Gardner EA, Johnston JA, et al. Double-blind comparison of bupropion and fluoxetine in depressed outpatients. *J Clin Psychiatry*. 1991 Aug;52(8):329-35. Epub: 1991/08/01. PMID: 1907963.
37. Fontaine R, Ontiveros A, Elie R, et al. A double-blind comparison of nefazodone, imipramine, and placebo in major depression. *J Clin Psychiatry*. 1994;55(6):234-41. PMID: 1995-02824-001. PMID: 8071277. Partial author list. First Author & Affiliation: Fontaine, R.
38. Frank E, Cassano GB, Rucci P, et al. Predictors and moderators of time to remission of major depression with interpersonal psychotherapy and SSRI pharmacotherapy. *Psychol Med*. 2011 Jan;41(1):151-62. Epub: 2010/04/13. PMID: 20380782.
39. Gastpar M, Singer A, Zeller K. Efficacy and tolerability of hypericum extract STW3 in long-term treatment with a once-daily dosage in comparison with sertraline. *Pharmacopsychiatry*. 2005 Mar;38(2):78-86. PMID: 15744631.
40. Gastpar M, Singer A, Zeller K. Comparative efficacy and safety of a once-daily dosage of hypericum extract STW3-VI and citalopram in patients with moderate depression: a double-blind, randomised, multicentre, placebo-controlled study. *Pharmacopsychiatry*. 2006 Mar;39(2):66-75. Epub: 2006/03/24. PMID: 16555167.
41. GmbH WP. A double-blind, placebo-controlled, comparative efficacy study of venlafaxine and sertraline in producing remission in outpatients with major depressive disorder: study 0600C1-402-US-CA-CSR-48579
42. GmbH WP. A double-blind, placebo-controlled, comparative study of extended release formulation of venlafaxine and imipramine on the time of onset of antidepressant response in patients with severe major depression: study 0600B1-384-US-EU-CA-CSR-41642.
43. GmbH WP. Randomized, double-blind comparison of venlafaxine (WY-45, 030), imipramine, and placebo capsules in outpatients with major depression: study 600A-303-US-303-EXT-GMR-20448.
44. GmbH WP. Randomized, double-blind comparison of venlafaxine (WY-45, 030), imipramine, and placebo capsules in outpatients with major depression: study 600A-303-US-303-EXT-GMR-20448.
45. GmbH WP. A randomized double-blind comparison of venlafaxine XR and paroxetine in outpatients with moderate to severe major depression: study 0600-428-IT-SDC-3993.
46. Golden RN, Nemeroff CB, McSorley P, et al. Efficacy and tolerability of controlled-release and immediate-release paroxetine in the treatment of depression. *J Clin Psychiatry*. 2002 Jul;63(7):577-84. Epub: 2002/07/30. PMID: 12143913.
47. Goldstein DJ, Lu Y, Detke MJ, et al. Duloxetine in the treatment of depression: a double-blind placebo-controlled comparison with paroxetine. *J Clin Psychopharmacol*. 2004 Aug;24(4):389-99. Epub: 2004/07/03. PMID: 15232330.
48. Goldstein DJ, Mallinckrodt C, Lu Y, et al. Duloxetine in the treatment of major depressive disorder: a double-blind clinical trial. *J Clin Psychiatry*. 2002 Mar;63(3):225-31. Epub: 2002/04/03. PMID: 11926722.
49. Hansgen KD, Vesper J, Ploch M. Multicenter double-blind study examining the antidepressant effectiveness of the hypericum extract LI 160. *J Geriatr Psychiatry Neurol*. 1994 Oct;7 Suppl 1:S15-8. Epub: 1994/10/01. PMID: 7857501.

50. Hegerl U, Hautzinger M, Mergl R, et al. Effects of pharmacotherapy and psychotherapy in depressed primary-care patients: a randomized, controlled trial including a patients' choice arm. *Int J Neuropsychopharmacol*. 2010;13(1):31-44. PMID: 19341510.
51. Henigsberg N, Mahableshwarkar AR, Jacobsen P, et al. A randomized, double-blind, placebo-controlled 8-week trial of the efficacy and tolerability of multiple doses of Lu AA21004 in adults with major depressive disorder. *J Clin Psychiatry*. 2012 Jul;73(7):953-9. Epub: 2012/08/21. PMID: 22901346.
52. Hicks JA, Argyropoulos S, Rich AS, et al. Randomised controlled study of sleep after nefazodone or paroxetine treatment in out-patients with depression. *Br J Psychiatry*. 2002;180(6):528-35. PMID: 2002-13724-013. First Author & Affiliation: Hicks, Jane A.
53. Higuchi T, Hong JP, Jung HY, et al. Paroxetine controlled-release formulation in the treatment of major depressive disorder: a randomized, double-blind, placebo-controlled study in Japan and Korea. *Psychiatry Clin Neurosci*. 2011 Dec;65(7):655-63. Epub: 2011/09/08. PMID: 21895859.
54. Hong CJ, Hu WH, Chen CC, et al. A double-blind, randomized, group-comparative study of the tolerability and efficacy of 6 weeks' treatment with mirtazapine or fluoxetine in depressed Chinese patients. *J Clin Psychiatry*. 2003;64(8):921-6.
55. Huang Y, Htut W, Li D, et al. Studies on the clinical observation and cerebral glucose metabolism in depression treated by electro-scalp acupuncture compared to fluoxetine. *Int J Clin Acupunct*. 2005;14(1):7-26. PMID: 0077160.
56. Iwata N, Tourian KA, Hwang E, et al. Efficacy and safety of desvenlafaxine 25 and 50/50% shaded blockmg/day in a randomized, placebo-controlled study of depressed outpatients. *J Psychiatr Pract*. 2013 Jan;19(1):5-14. Epub: 2013/01/22. PMID: 23334675.
57. Jazayeri S, Tehrani-Doost M, Keshavarz SA, et al. Comparison of therapeutic effects of omega-3 fatty acid eicosapentaenoic acid and fluoxetine, separately and in combination, in major depressive disorder. *Aust N Z J Psychiatry*. 2008 Mar;42(3):192-8. Epub: 2008/02/06. PMID: 18247193.
58. Kalb R, Trautmann-Sponsel RD, Kieser M. Efficacy and tolerability of hypericum extract WS 5572 versus placebo in mildly to moderately depressed patients. A randomized double-blind multicenter clinical trial. *Pharmacopsychiatry*. 2001;34(3):96-103. PMID: CN-00354554.
59. Kasper S, Olivieri L, Di Loreto G, et al. A comparative, randomised, double-blind study of trazodone prolonged-release and paroxetine in the treatment of patients with major depressive disorder. *Curr Med Res Opin*. 2005 Aug;21(8):1139-46. Epub: 2005/08/09. PMID: 16083521.
60. Kennedy SH, Konarski JZ, Segal ZV, et al. Differences in brain glucose metabolism between responders to CBT and venlafaxine in a 16-week randomized controlled trial. *Am J Psychiatry*. 2007 May;164(5):778-88. Epub: 2007/05/04. PMID: 17475737.
61. Khan A, Bose A, Alexopoulos GS, et al. Double-blind comparison of escitalopram and duloxetine in the acute treatment of major depressive disorder. *Clin Drug Investig*. 2007;27(7):481-92. Epub: 2007/06/15. PMID: 17563128.
62. Laakmann G, Schule C, Baghai T, et al. St. John's wort in mild to moderate depression: the relevance of hyperforin for the clinical efficacy. *Pharmacopsychiatry*. 1998 Jun;31 Suppl 1:54-9. Epub: 1998/07/31. PMID: 9684948.
63. Lee P, Shu L, Xu X, et al. Once-daily duloxetine 60 mg in the treatment of major depressive disorder: multicenter, double-blind, randomized, paroxetine-controlled, non-inferiority trial in China, Korea, Taiwan and Brazil. *Psychiatry Clin Neurosci*. 2007 Jun;61(3):295-307. Epub: 2007/05/03. PMID: 17472599.

64. Liebowitz MR, Tourian KA, Hwang E, et al. A double-blind, randomized, placebo-controlled study assessing the efficacy and tolerability of desvenlafaxine 10 and 50 mg/day in adult outpatients with major depressive disorder. *BMC Psychiatry*. 2013;13:94. Epub: 2013/03/23. PMID: 23517291.
65. Liebowitz MR, Yeung PP, Entsuah R. A randomized, double-blind, placebo-controlled trial of desvenlafaxine succinate in adult outpatients with major depressive disorder. *J Clin Psychiatry*. 2007 Nov;68(11):1663-72. Epub: 2007/12/07. PMID: 18052559.
66. Lydiard RB, Stahl SM, Hertzman M, et al. A double-blind, placebo-controlled study comparing the effects of sertraline versus amitriptyline in the treatment of major depression. *J Clin Psychiatry*. 1997 Nov;58(11):484-91. Epub: 1997/12/31. PMID: 9413414.
67. Mehtonen OP, Sogaard J, Roponen P, et al. Randomized, double-blind comparison of venlafaxine and sertraline in outpatients with major depressive disorder. Venlafaxine 631 Study Group. *J Clin Psychiatry*. 2000 Feb;61(2):95-100. Epub: 2000/03/25. PMID: 10732656.
68. Mischoulon D, Price LH, Carpenter LL, et al. A double-blind, randomized, placebo-controlled clinical trial of S-adenosyl-L-methionine (S-AMe) versus escitalopram in major depressive disorder. *J Clin Psychiatry*. 2014 Dec 24. Epub: 2014/02/07. PMID: 24500245.
69. Montgomery SA, Mansuy L, Ruth A, et al. Efficacy and safety of levomilnacipran sustained release in moderate to severe major depressive disorder: a randomized, double-blind, placebo-controlled, proof-of-concept study. *J Clin Psychiatry*. 2013 Apr;74(4):363-9. Epub: 2013/05/10. PMID: 23656841.
70. Munizza C, Olivieri L, Di Loreto G, et al. A comparative, randomized, double-blind study of trazodone prolonged-release and sertraline in the treatment of major depressive disorder. *Curr Med Res Opin*. 2006 Sep;22(9):1703-13. Epub: 2006/09/14. PMID: 16968574.
71. Nemeroff CB, Thase ME. A double-blind, placebo-controlled comparison of venlafaxine and fluoxetine treatment in depressed outpatients. *J Psychiatr Res*. 2007;41(3-4):351-9. PMID: 2006-22968-021. PMID: 16165158. First Author & Affiliation: Nemeroff, Charles B.
72. Nierenberg AA, Greist JH, Mallinckrodt CH, et al. Duloxetine versus escitalopram and placebo in the treatment of patients with major depressive disorder: onset of antidepressant action, a non-inferiority study. *Curr Med Res Opin*. 2007 Feb;23(2):401-16. Epub: 2007/02/10. PMID: 17288694.
73. Oakes TM, Myers AL, Marangell LB, et al. Assessment of depressive symptoms and functional outcomes in patients with major depressive disorder treated with duloxetine versus placebo: primary outcomes from two trials conducted under the same protocol. *Hum Psychopharmacol*. 2012 Jan;27(1):47-56. Epub: 2012/01/14. PMID: 22241678.
74. Olie JP, Gunn KP, Katz E. A double-blind placebo-controlled multicentre study of sertraline in the acute and continuation treatment of major depression. *Eur Psychiatry*. 1997;12(1):34-41. PMID: CN-00189012.
75. Perahia DG, Wang F, Mallinckrodt CH, et al. Duloxetine in the treatment of major depressive disorder: a placebo- and paroxetine-controlled trial. *Eur Psychiatry*. 2006 Sep;21(6):367-78. Epub: 2006/05/16. PMID: 16697153.
76. Philipp M, Kohnen R, Hiller KO. Hypericum extract versus imipramine or placebo in patients with moderate depression: randomised multicentre study of treatment for eight weeks. *BMJ*. 1999 Dec 11;319(7224):1534-8. Epub: 1999/12/11. PMID: 10591711.
77. Qu SS, Huang Y, Zhang ZJ, et al. A 6-week randomized controlled trial with 4-week follow-up of acupuncture combined with paroxetine in patients with major depressive disorder. *J Psychiatr Res*. 2013 Jun;47(6):726-32. Epub: 2013/03/19. PMID: 23498306.

78. Quah-Smith I, Smith C, Crawford JD, et al. Laser acupuncture for depression: a randomised double blind controlled trial using low intensity laser intervention. *J Affect Disord.* 2013 Jun;148(2-3):179-87. Epub: 2013/01/23. PMID: 23337655.
79. Reimherr FW, Chouinard G, Cohn CK, et al. Antidepressant efficacy of sertraline: a double-blind, placebo- and amitriptyline-controlled, multicenter comparison study in outpatients with major depression. *J Clin Psychiatry.* 1990 Dec;51 Suppl B:18-27. Epub: 1990/12/01. PMID: 2258378.
80. Rickels K, Amsterdam J, Clary C, et al. The efficacy and safety of paroxetine compared with placebo in outpatients with major depression. *J Clin Psychiatry.* 1992 Feb;53 Suppl:30-2. Epub: 1992/02/01. PMID: 1531820.
81. Rickels K, Schweizer E, Clary C, et al. Nefazodone and imipramine in major depression: a placebo-controlled trial. *Br J Psychiatry.* 1994 Jun;164(6):802-5. Epub: 1994/06/01. PMID: 7952987.
82. Rudolph RL, Fabre LF, Feighner JP, et al. A randomized, placebo-controlled, dose-response trial of venlafaxine hydrochloride in the treatment of major depression. *J Clin Psychiatry.* 1998 Mar;59(3):116-22. Epub: 1998/04/16. PMID: 9541154.
83. Rudolph RL, Feiger AD. A double-blind, randomized, placebo-controlled trial of once-daily venlafaxine extended release (XR) and fluoxetine for the treatment of depression. *J Affect Disord.* 1999 Dec;56(2-3):171-81. Epub: 2000/03/04. PMID: 10701474.
84. Rush AJ, Trivedi MH, Carmody TJ, et al. Response in relation to baseline anxiety levels in major depressive disorder treated with bupropion sustained release or sertraline. *Neuropsychopharmacology.* 2001 Jul;25(1):131-8. Epub: 2001/05/30. PMID: 11377926.
85. Schrader E. Equivalence of St John's wort extract (Ze 117) and fluoxetine: a randomized, controlled study in mild-moderate depression. *Int Clin Psychopharmacol.* 2000 Mar;15(2):61-8. PMID: 10759336.
86. Septien-Velez L, Pitrosky B, Padmanabhan SK, et al. A randomized, double-blind, placebo-controlled trial of desvenlafaxine succinate in the treatment of major depressive disorder. *Int Clin Psychopharmacol.* 2007 Nov;22(6):338-47. Epub: 2007/10/06. PMID: 17917552.
87. Shahidi M, Mojtahed A, Modabbernia A, et al. Laughter Yoga versus group exercise program in elderly depressed women: A randomized controlled trial. *Int J Geriatr Psychiatry.* 2011;26(3):322-7. PMID: 2011-05494-014. PMID: 20848578. First Author & Affiliation: Shahidi, Mahvash.
88. Sheehan DV, Croft HA, Gossen ER, et al. Extended-release Trazodone in Major Depressive Disorder: A Randomized, Double-blind, Placebo-controlled Study. *Psychiatry (Edmont).* 2009 May;6(5):20-33. Epub: 2009/09/03. PMID: 19724732.
89. Shelton RC, Haman KL, Rapaport MH, et al. A randomized, double-blind, active-control study of sertraline versus venlafaxine XR in major depressive disorder. *J Clin Psychiatry.* 2006 Nov;67(11):1674-81. Epub: 2007/01/02. PMID: 17196045.
90. Silverstone PH, Ravindran A. Once-daily venlafaxine extended release (XR) compared with fluoxetine in outpatients with depression and anxiety. Venlafaxine XR 360 Study Group. *J Clin Psychiatry.* 1999 Jan;60(1):22-8. Epub: 1999/03/13. PMID: 10074873.
91. Sir A, D'Souza RF, Uguz S, et al. Randomized trial of sertraline versus venlafaxine XR in major depression: efficacy and discontinuation symptoms. *J Clin Psychiatry.* 2005 Oct;66(10):1312-20. Epub: 2005/11/02. PMID: 16259546.
92. Smith WT, Glaudin V. A placebo-controlled trial of paroxetine in the treatment of major depression. *J Clin Psychiatry.* 1992 Feb;53 Suppl:36-9. Epub: 1992/02/01. PMID: 1531822.
93. Smith WT, Glaudin V, Panagides J, et al. Mirtazapine vs. amitriptyline vs. placebo in the treatment of major depressive disorder. *Psychopharmacol Bull.* 1990;26(2):191-6. Epub: 1990/01/01. PMID: 2236455.

94. Soares CN, Thase ME, Clayton A, et al. Desvenlafaxine and escitalopram for the treatment of postmenopausal women with major depressive disorder. *Menopause*. 2010 Jul;17(4):700-11. Epub: 2010/06/12. PMID: 20539246.
95. Stevinson C. Negative result for St John's wort in major depression. *Focus on Alternative and Complementary Therapies*. 2001 Sep;6(3):196-7. PMID: 0029317.
96. Sun H, Zhao H, Ma C, et al. Effects of electroacupuncture on depression and the production of glial cell line-derived neurotrophic factor compared with fluoxetine: a randomized controlled pilot study. *J Altern Complement Med*. 2013 Sep;19(9):733-9. PMID: 23647408.
97. Szegedi A, Kohlen R, Dienel A, et al. Acute treatment of moderate to severe depression with hypericum extract WS 5570 (St John's wort): randomised controlled double blind non-inferiority trial versus paroxetine. *BMJ*. 2005 Mar 5;330(7490):503. Epub: 2005/02/15. PMID: 15708844.
98. Thase ME. Efficacy and tolerability of once-daily venlafaxine extended release (XR) in outpatients with major depression. The Venlafaxine XR 209 Study Group. *J Clin Psychiatry*. 1997 Sep;58(9):393-8. Epub: 1997/10/29. PMID: 9378690.
99. Tourian KA, Padmanabhan SK, Groark J, et al. Desvenlafaxine 50 and 100 mg/d in the treatment of major depressive disorder: an 8-week, phase III, multicenter, randomized, double-blind, placebo-controlled, parallel-group trial and a post hoc pooled analysis of three studies. *Clin Ther*. 2009 Jun;31 Pt 1:1405-23. Epub: 2009/08/25. PMID: 19698901.
100. Uebelhack R, Gruenwald J, Graubaum HJ, et al. Efficacy and tolerability of Hypericum extract STW 3-VI in patients with moderate depression: a double-blind, randomized, placebo-controlled clinical trial. *Adv Ther*. 2004 Jul-Aug;21(4):265-75. Epub: 2004/12/21. PMID: 15605620.
101. van Moffaert M, de Wilde J, Vereecken A, et al. Mirtazapine is more effective than trazodone: a double-blind controlled study in hospitalized patients with major depression. *Int Clin Psychopharmacol*. 1995 Mar;10(1):3-9. Epub: 1995/03/01. PMID: 7622801.
102. Ventura D, Armstrong EP, Skrepnek GH, et al. Escitalopram versus sertraline in the treatment of major depressive disorder: a randomized clinical trial. *Curr Med Res Opin*. 2007 Feb;23(2):245-50. Epub: 2007/02/10. PMID: 17288677.
103. Wade A, Crawford GM, Angus M, et al. A randomized, double-blind, 24-week study comparing the efficacy and tolerability of mirtazapine and paroxetine in depressed patients in primary care. *Int Clin Psychopharmacol*. 2003 May;18(3):133-41. Epub: 2003/04/19. PMID: 12702891.
104. Wade A, Gembert K, Florea I. A comparative study of the efficacy of acute and continuation treatment with escitalopram versus duloxetine in patients with major depressive disorder. *Curr Med Res Opin*. 2007 Jul;23(7):1605-14. Epub: 2007/06/15. PMID: 17559755.